

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau


INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/82, A01H 5/00	A2	(11) International Publication Number: WO 00/50613 (43) International Publication Date: 31 August 2000 (31.08.00)
(21) International Application Number: PCT/IL00/00110 (22) International Filing Date: 22 February 2000 (22.02.00) (30) Priority Data: 60/121,239 22 February 1999 (22.02.99) US (71) Applicant (for all designated States except US): YISSUM RESEARCH AND DEVELOPMENT COMPANY OF THE HEBREW UNIVERSITY OF JERUSALEM [IL/IL]; Jabotinsky Street 46, 91042 Jerusalem (IL). (72) Inventors; and (75) Inventors/Applicants (for US only): VAINSTEIN, Alexander [IL/IL]; Hanassi Harishron Street 42/6, 76303 Rehovot (IL). ZUKER, Amir [IL/IL]; Negba Street 32/18, 74040 Nes Ziona (IL). OVADIS, Marianna [IL/IL]; Hershenson Street 76/28, 76484 Rehovot (IL). (74) Agent: REINHOLD COHN AND PARTNERS; P.O. Box 4060, 61040 Tel Aviv (IL).	(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published Without international search report and to be republished upon receipt of that report.	

(54) Title: TRANSGENIC PLANTS AND METHOD FOR TRANSFORMING CARNATIONS

(57) Abstract

A method of transforming a carnation (*Dianthus l.*) plant genome with a DNA molecule. The method comprises (a) preparing stem explants from carnation cuttings; (b) wounding the explants by microprojectile bombardment; (c) cocultivating the wounded explants with *Agrobacterium* comprising the DNA molecule under defined conditions of exposure to dark followed by light; (d) excising shoots from the cultivated wounded explants and removing the leaves from the shoots; and (e) culturing the leaves to obtain transgenic shoots transformed with the DNA molecule. Also disclosed are a *rolC*-transgenic carnation with improved agronomic traits and enhancement of flower fragrance by antisense suppression of the flavonoid gene *fhl*.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IL 00/00110

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0486233 A		DE 69111385 D	24-08-1995
		DE 69111385 T	04-01-1996
		DK 486233 T	11-09-1995
		ES 2077181 T	16-11-1995
		GR 3017619 T	31-01-1996
		HU 60783 A	28-10-1992
		JP 5308960 A	22-11-1993
		NZ 239977 A	26-08-1993
WO 9318142 A	16-09-1993	AU 668548 B	09-05-1996
		AU 3786493 A	05-10-1993
		BG 99033 A	29-09-1995
		BR 9306047 A	18-11-1997
		CA 2131704 A	16-09-1993
		EP 0630403 A	28-12-1994
		HU 69991 A	28-09-1995
		JP 7506963 T	03-08-1995
		NZ 251069 A	26-01-1996
		US 5733759 A	31-03-1998
WO 9715584 A	01-05-1997	AU 7663296 A	15-05-1997
		EP 0858507 A	19-08-1998
WO 9850570 A	12-11-1998	AU 7377298 A	27-11-1998
WO 9721816 A	19-06-1997	AU 1106297 A	03-07-1997

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IL 00/00110

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9217056 A	15-10-1992	EP 0582603 A	16-02-1994
		US 5589613 A	31-12-1996
WO 9639827 A	19-12-1996	AU 4648996 A	30-12-1996
		BR 9508561 A	19-05-1998
		CA 2196972 A	19-12-1996
		EP 0776159 A	04-06-1997
		NZ 301140 A	23-12-1998
		US 5792467 A	11-08-1998
		ZA 9600006 A	27-08-1996
WO 9620595 A	11-07-1996	AU 4744296 A	24-07-1996
		BR 9510180 A	14-10-1997
		CA 2208753 A	11-07-1996
		EP 0800345 A	15-10-1997
		EP 1000543 A	17-05-2000
		JP 11501295 T	02-02-1999
		PL 321179 A	24-11-1997
		ZA 9600005 A	27-08-1996
WO 9735471 A	02-10-1997	AU 3202597 A	17-10-1997
		CA 2250182 A	02-10-1997
		EP 0889691 A	13-01-1999
WO 9636716 A	21-11-1996	AU 699874 B	17-12-1998
		AU 5639696 A	29-11-1996
		CA 2202668 A	21-11-1996
		EP 0873410 A	28-10-1998
		HU 9802555 A	01-02-1999
		JP 11505116 T	18-05-1999
		NZ 307119 A	29-04-1999
		US 6080920 A	27-06-2000
WO 9937794 A	29-07-1999	AU 2424199 A	09-08-1999
WO 9428140 A	08-12-1994	AU 672308 B	26-09-1996
		AU 6789594 A	20-12-1994
		CA 2163220 A	08-12-1994
		CN 1127015 A	17-07-1996
		EP 0703982 A	03-04-1996
		JP 8511683 T	10-12-1996
		NZ 266401 A	25-09-1996
		PL 311691 A	04-03-1996
		SG 45175 A	16-01-1998
		US 5948955 A	07-09-1999
WO 9506741 A	09-03-1995	FR 2709496 A	10-03-1995
		AU 695398 B	13-08-1998
		AU 7539694 A	22-03-1995
		EP 0716706 A	19-06-1996
		HU 74664 A	28-01-1997
		JP 9509302 T	22-09-1997
		SK 25896 A	04-09-1996
EP 0486233 A	20-05-1992	AT 125301 T	15-08-1995
		AU 643846 B	25-11-1993
		AU 8771391 A	21-05-1992
		CA 2053473 A	15-05-1992

transgenic carnation plants with novel characteristics. IX International Congress on Plant Tissue and Cell Culture, Jerusalem, Israel (1998)

7. Fladung, M., K. Grossmann, and M.R. Ahuja. 1997. Alterations in hormonal and developmental characteristics in transgenic *Populus* conditioned by the *rolC* gene from *Agrobacterium rhizogenes*. J. Plant Physiol. 150:420-427
8. Scorza, R., T.W. Zimmerman, J.M. Cordts, K.J. Footen, and M. Ravelonandro. 1994. Horticultural characteristics of transgenic tobacco expressing *rolC* gene from *Agrobacterium rhizogenes*. J. Amer. Soc. Hort. Sci. 119:1091-1098
9. Nilsson, O. and O. Olsson. 1997. Getting to the root: the role of the *Agrobacterium rhizogenes rol* genes in the formation of hairy roots. Physiol. Plant. 100:463-473
10. Souq, F., P. Coutos-Thevenot, H. Yean, G. Delbard, Y. Maziere, J.P. Barbe, and M. Boulay. 1986. Genetic transformation of roses, 2 examples: one on morphogenesis, the other on anthocyanin biosynthetic pathway. In: Morisot A, Ricci P (eds) Second International Symposium on Roses. Acta Hort. 424:381-388
11. Pelletier, M.K. & Shirley, B.W. Analysis of flavanone 3-hydroxylase in arabidopsis seedlings. *Plant Physiol.* 111, 339-345 (1996)
12. Ahroni A: Developing efficient regeneration and transformation methods for carnation and gypsophila. M.Sc. thesis (The Hebrew University of Jerusalem, Israel) (1996)
13. Tzfira, T., C.S. Jensen, W. Wang, A. Zuker, B. Vinocur, A. Altman, and A. Vainstein. 1997. Transgenic *Populus tremula*: a step-by-step protocol for its *Agrobacterium*-mediated transformation. Plant Mol. Biol. Rep. 15:219-235

25 The carnation (*Dianthus* L.), one of the world's major cut-flower crops, is the commercial leader in terms of number of stems sold worldwide. Different carnation types have been developed as a result of crossing and they can be divided into two main groups, standard (midi and mignon carnation) and spray (micro and diantini carnation); a minor group is formed by the pot carnations.

- 1 -

TRANSGENIC PLANTS AND METHOD FOR TRANSFORMING CARNATIONS

FIELD OF THE INVENTION

This invention relates to a method for transforming carnation plants and novel transgenic plants with improved traits.

BACKGROUND OF THE INVENTION

5 The following references may contribute to the understanding of the invention, and are referred to by number in the specification:

1. Zuker A, Tzfira T, Vainstein A: Genetic engineering for cut-flower improvement. *Biotech Adv* 16: 33-79 (1998)
2. van Altvorst AC, Rikesen T, Koehorst H, Dons JJM: Transgenic carnations
10 obtained by *Agrobacterium tumefaciens*-mediated of leaf explants. *Transgenic Res* 4: 105-113 (1995)
3. Zuker A, Chang P-FL, Ahroni A, Cheah K, Woodson WR, Bressan RA, Watad AA, Hasegawa PM, Vainstein A: Transformation of carnation by microprojectile bombardment. *Sci Hort* 64: 177-185 (1995)
- 15 4. Firoozabady E, Moy Y, Tucker W, Robinson K, Gutterson N: Efficient transformation and regeneration of carnation cultivars using *Agrobacterium*. *Molecular Breeding* 1: 283-293 (1995)
5. Lu CY, Nugent G, Wardley-Richardson T, Chandler SF, Young R, Dalling MJ: *Agrobacterium*-mediated transformation of carnation (*Dianthus caryophyllus*
20 L.). *Bio/Tech* 9: 864-868 (1991)
6. Zuker, A., et al: Application of an integrative system based on microprojectile bombardment and *Agrobacterium tumefaciens* to generate

- 4 -

microprojectile bombardment and *Agrobacterium*-mediated transformation under a light regime and a two-step selection cycle has recently appeared (6).

New traits in cut flowers include not only yield improvement and resistance to insects or disease; they also consist of new colors and novel plant morphology. In fact, the latter are of great importance for the cut-flower market, where plant architecture and flower color are the main features determining consumer interest. Nevertheless, agronomic traits, such as performance, remain highly important for breeders and growers as extensively growing and better performing plants can lower the time and cost required for growth and breeding.

Among the different genes affecting plant morphology, e.g. homeotic, *Agrobacterium*, phytochrome and gibberellin genes, the *Agrobacterium rhizogenes* *rol* genes have been the most widely and successfully employed (7,8). Although its precise mode of action is still unknown (9), *rolC* has attracted the most attention through its expression in transgenic plants—either under its own promoter or under the control of a cauliflower mosaic virus (CaMV) 35S promoter, leading to a series of morphological alterations. These include reduced apical dominance, altered leaf morphology, reduced seed production, reduced internode length, male sterility, small flowers and early flowering, bushy and compact phenotype, and even stem fasciation.

Nevertheless, studies on the *rolC* gene and its effects on plant development have been performed mostly in model herbaceous plants, e.g. tobacco, potato, and tomato, or forest trees. *rolC* studies in cut flowers are rather limited—the only report being on a woody ornamental, rose, in the proceedings of a symposium (10). These transgenic rose plants (*Rosa hybrida* cv. Madame G. Delbard), expressing the *rolC* gene under its native promoter, exhibited an array of phenotypic alterations, most of which were highly disadvantageous horticulturally. They included a dramatically reduced root system, wrinkled leaves, high sensitivity to diseases and dying out of shoots.

With respect to color, anthocyanins, carotenoids and betalain are the main flower pigments, of which the most studied are the anthocyanins. The great interest

- 3 -

Market demand for flowers with improved traits—such as new colors, new shapes, better fragrance and longer vase life, drives breeders to create new and more attractive varieties every year. However, carnation is a vegetatively propagated crop and is very detrimentally affected by inbreeding. Hence, controlled breeding is rather complicated and limited due to the fact that selection of a desired trait in the siblings is performed on the genetic background of their two parents, and due to the very high genetic variability among offspring. Furthermore, crosses within and between related species is limited by a rather small available gene pool for new traits.

Biotechnological techniques such as genetic engineering could be a useful alternative/addition to currently used classical breeding methods for the production of novel plant varieties (1).

One of the prerequisites to generating transgenic plants is the availability of a regeneration system. For carnation, numerous procedures leading to efficient adventitious regeneration from a number of explants, including stem segments, leaves and petals, have been published (1). However, success in harnessing these regeneration systems to efficiently generate transgenes has been rather limited. The *Agrobacterium*-mediated transformation procedure described by van Altvorst et al. (2) and the microprojectile bombardment-mediated procedure (3) were both of relatively low efficiency.

A more recent, alternative transformation procedure was developed by DNAP Inc. (4). Although a high number of transgenes could be generated from the three varieties used, the procedure was extremely time-consuming and cumbersome because it relied on vitrified leaves as the primary explants and these can take 4 to 6 months of tissue culture to generate. The only efficient procedure reported to date, also used to generate transgenes with improved vase life, was developed by another biotech company, Calgene Pacific Ltd. (5). However extensive efforts by numerous groups to generate carnation transgenes with this experimental protocol have been unsuccessful. A preliminary report of an integrative system based on

- 6 -

- (c) cocultivating the wounded explants with *Agrobacterium* comprising the DNA molecule under conditions of exposure to dark followed by light;
- (d) excising shoots from the cultivated wounded explants and removing the leaves from the shoots; and
- 5 (e) culturing the leaves to obtain transgenic shoots transformed with the DNA molecule.

The method of the invention combines three experimental conditions to produce an efficient carnation transformation procedure: (1) the use of microprojectile bombardment to wound stem explants, (2) a specific dark/light 10 regime to optimize transformation, and (3) an optimized two-cycle selection procedure.

Stem explants may be prepared from (stem) cuttings with e.g., four to ten fully mature leaves (not counting the apical leaves which were not fully expanded), harvested from greenhouse-grown plants and stored for up to 1 month at 4°C. The 15 leaves and shoot apices may be removed and primary nodes (two to five) isolated.

Each bombardment typically consists of 50-200 μ l of tungsten or gold particles in an aqueous suspension (10-20mg/ml). Explants are generally placed on medium solidified with 1% agar and bombarded 1-2 times. Preferred particle acceleration parameters are: a bombardment pressure of 1300-2000 psi and a 20 distance of 3-12 cm from the launching plate to the tissue.

Inoculated stem explants are cultured in an upright position for a period of time in the dark followed by a period of light. In a preferred embodiment, the explants are cocultivated for 2-4 days in the dark and then 1-4 days in the light. In a most preferred embodiment, the cultivation is carried out for 3 days in the dark 25 followed by 2 days in the light.

For the first selection cycle during shoot regeneration from the stem explants, the basic medium was preferably supplemented with 0.04-1 mg/l NAA and 0.6-3 mg/l TDZ (SI-T1). After ca. 14-28 days, clusters of regenerated adventitious shoots are excised from the primary stem explants. Leaves from all of 30 the shoots of each independent cluster are pulled off and cultured in medium

- 5 -

in the latter, which derive from the general phenylpropanoid pathway, results from their wide distribution in many plants and microorganisms and their role in color determination. Furthermore, detailed biochemical and genetic analyses of anthocyanin production/accumulation has brought about the development of two
5 main strategies for altering flower color: introducing a foreign gene(s) to allow new branching in the anthocyanin-biosynthesis pathway, and up/down-regulation of this pathway's native genes' expression.

Several genes from the anthocyanin biosynthetic pathway have been used to manipulate flower color in numerous plants. Chalcone synthase (*chs*), chalcone
10 flavanone isomerase (*chi*) and flavanone 3-hydroxylase (*fht*) are the first three genes encoding the early, non-branched segment of the flavonoid biosynthetic pathway (Fig. 17). Although FHT is a key enzyme in this pathway (11) and its suppression, as opposed to that of CHS, should not block the production of the essential phytoalexins, isoflavonoids, it has never been harnessed for the genetic
15 manipulation of flower color.

With respect to flower fragrance, mainly dominated by terpenoids, fatty acid derivatives and benzenoid compounds (phenylpropanoid derivatives), much less is known. To date, only a few genes involved in fragrance production have been characterized.

20 SUMMARY OF THE INVENTION

It is an object of the present invention to provide a method for transforming carnation plants.

It is a further object of the invention to provide transgenic carnation plants with desirable traits.

25 In one aspect of the invention, there is provided a method of transforming a carnation (*Dianthus L.*) plant genome with a DNA molecule comprising:

- (a) preparing stem explants from carnation cuttings;
- (b) wounding the explants by microprojectile bombardment;

- 8 -

In a preferred embodiment of this aspect of the invention, the expression of the *fht* gene is blocked by an antisense molecule.

In a most preferred embodiment of this aspect of the invention, the antisense molecule comprises the sequence of Fig. 13.

5 In a preferred embodiment, the transgenic carnation plant is produced by the method of the first aspect of the invention.

In this aspect of the invention, expression of a gene encoding flavanone 3-hydroxylase (*fht*), a key step in the anthocyanin pathway is blocked by antisense suppression in carnation. Transgenic carnation plants may exhibit flower color
10 modifications ranging from attenuation to complete loss of their original color. Hence, carnation flowers in a variety of colors can be generated.

Approach to olfactory enhancement of natural fragrance (flower fragrance in particular) via diverting metabolic flow from anthocyanins to the production of benzoic acid derivatives (fragrance compounds) is demonstrated in carnation. In a
15 preferred embodiment, transgenic carnation plants with a blocked flavonoid biosynthetic pathway (expression of the *fht* gene) may be generated using the above mentioned transformation procedure of the first aspect of the invention. Flowers of these carnation transgenes are much more fragrant than control plants.

In a fourth aspect of the invention, there is provided a method for
20 controlling the fragrance of a plant comprising modulating gene expression in the anthocyanin-biosynthetic pathway of the plant.

In a preferred embodiment, the modulation comprises suppressing the expression of one or more genes in the pathway. An example of such a gene is the *fht* gene. In a most preferred embodiment, the suppression is by an antisense
25 molecule.

DETAILED DESCRIPTION OF THE DRAWINGS

In order to understand the invention and to see how it may be carried out in practice, a preferred embodiment will now be described, by way of non-limiting example only, with reference to the accompanying drawings, in which:

- 7 -

supplemented with 0.04-1 mg/l NAA and 0.5-2 mg/l BAP (SI-B1) for adventitious shoot regeneration and selection of transgenes (second selection cycle). After 10 to 20 days, new adventitious transgenic shoots emerge from the leaf basal area.

In a second aspect of the invention, there is provided a transgenic carnation plant transformed with the *Agrobacterium rhizogenes rolC* gene.

In a preferred embodiment of this aspect of the invention, the *rolC* gene is under the control of a cauliflower mosaic virus (CaMV) 35S promoter.

In a further preferred embodiment of this aspect of the invention, the transgenic carnation plant is produced by the method of the first aspect of the invention.

In a most preferred embodiment of this aspect of the invention, the gene used to transform the carnation comprises the sequence shown in Fig. 6.

rolC-transgenic carnation plants (or "*rolC* transgenes") according to the invention generally exhibit increased axillary bud breakage and development when grown under standard commercial greenhouse conditions. *rolC* transgenes often generate up to 40% more stem cuttings per mother plant than controls. Stem cuttings from *rolC* transgenes usually exhibit better rooting ability, with up to five times higher root dry weight than controls. The improved rooting of *rolC*-transgenic cuttings is also apparent when commercial rooting powder is used. Analyses of plants during the flowering season reveal that *rolC* transgenes generally yield up to three times more flowering stems than control plants.

Improved rooting, for example, can lower the labor required to produce rooted stem cuttings. The importance of these traits is even more pronounced for commercial varieties that exhibit very low rooting percentages despite the application of rooting powder. With respect to the yield characteristics, e.g. number of stem cuttings and flowering stems per mother plant, these are obviously of major importance to carnation propagators and growers.

In a third aspect of the invention, there is provided a transgenic carnation plant wherein expression of the flavanone 3-hydroxylase (*fht*) gene is blocked.

- 10 -

plants (C); and plasmid pCGN7001 (P). Following *nptII*-PCR amplification and gel electrophoresis, the products were hybridized with *nptII* probe.

Fig. 6 shows the DNA sequence of the *Agrobacterium rhizogenes rolC* gene.

Fig. 7 illustrates a schematic illustration of a typical carnation stem cutting. Each cutting is composed of three to five internodes (I to V). Each node carries two opposite leaves and two axillary buds which can develop into side shoots. The developmental stage of these side shoots was graded according to the number of open pairs of leaves on them (0, 1 or 2).

Figs. 8A and 8B show a molecular analysis of three independent *rolC*-transgenic (R-1, R-2 and R-3) and control *uidA*-transgenic (C) cv. White Sim carnation plants. (A) Southern blot analysis of *HindIII*-digested total DNA. (B) Northern blot analysis of total RNA. Both blots were hybridized with *rolC* probe.

Fig. 9 illustrates a distribution of stem-cutting types in *rolC*-transgenic (R-1, R-2 and R-3) and control (C) carnation plants. Stem cuttings were classified as III, IV or V according to the number of internodes they carried, as depicted in Fig. 7.

Fig. 10 illustrates adventitious root dry weight and rooting percentages in stem cuttings of *rolC*-transgenic lines (R-1, R-2 and R-3) and control (C) carnation plants after 30 days on rooting tables, as detailed in Materials and Methods of Example II. Bars represent an average of 15 stem cuttings per line; SE of the mean is indicated. The numbers above the bars represent the rooting percentages, i.e. percent of cuttings developing roots out of total number of cuttings.

Figs. 11a, 11b, 11c and 11d show morphological alterations in *rolC*-transgenic carnation plants: (11a) Rooted stem cuttings of control (left) and *rolC*-transgenic line R-1 (right). (11b) Flowers of *rolC*-transgenic line R-2 (top) and control (bottom) plants. (11c) Adventitious root formation in line R-1 (left) and control (right) plants. (11d) Flowering stems of *rolC*-transgenic lines and control plants (from left to right: R-1, R-2, R-3, control).

Fig. 12 shows the cDNA sequence of *fht* from carnation.

Fig. 13 shows the sequence of the antisense fragment to the *fht* cDNA.

- 9 -

Fig. 1 shows the effect of wounding and cocultivation on the efficiency of transient transformation of cv. White Sim stem explants. Explants were wounded by bombardment (+) or just poking/scratching and cocultivated with EHA105/pKIWI105 for 5 days under constant light or in the dark. GUS expression is presented as the number of blue spots per stem explant. S.E. of the mean ($p = 0.05$) is indicated ($n = 5$).

Fig. 2 illustrates transient transformation frequencies of different carnation cultivars. Following bombardment, stem explants were cocultivated with EHA105/pKIWI105 for 3 days in the dark and 2 days under constant light as described in Materials and Methods. GUS expression is presented as the number of blue spots per stem explant. S.E. of the mean ($p = 0.05$) is indicated ($n = 5$).

Figs. 3A-H illustrate transformation and regeneration of transgenic carnation plants. (A) GUS expression on the cut surface of a stem explant 14 days after inoculation. (B) Shoot regeneration from a stem explant. (C) The chimeric pattern of GUS expression following the first selection cycle. (D) Second selection cycle of adventitious shoots. Shoots developed from the leaf area which showed resistance to kanamycin, as reflected by its green color. The white, non-resistant area remained non-regenerative (arrow). (E,F) Solid, non-chimeric GUS expression in adventitious shoots regenerated from leaves following the second selection cycle. (G) Transgenic flowers. (H) GUS-expressing seedlings (T_1) obtained from crosses of transgenic cv. White Sim with non-transformed (b.l. 13261/2/3) male plants.

Fig. 4 shows southern blot analysis of DNA from four independent transgenic (1-4) and one non-transformed (C) cv. White Sim carnation plants. Total DNA (10 μ g) was digested with *Eco*RI (left) or *Hind*III (right) and hybridized with a *uidA* probe. (P) Plasmid pCGN7001 digested with *Eco*RI.

Fig. 5 shows a Southern blot of PCR analysis of offspring (T_1). DNA for *np11* PCR analyses was prepared from GUS-expressing kanamycin-resistant T_1 progeny plants (T_0 cv. White Sim x b.l. 13261); T_0 plants; control, non-transformed

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

I. METHOD OF CARNATION TRANSFORMATION

A. Materials and methods

1. Plant material

5 Carnation plants were grown under standard greenhouse conditions. The experiments described below were carried out using the carnation species *Dianthus caryophyllus* L. However, it is to be understood that this is only an exemplary species, and this aspect of the invention applies to all species of the *Dianthus* genus. Cultivars White Sim, Eilat, Darling, Visa and Lior were grown at the
10 Faculty of Agriculture (The Hebrew University of Jerusalem, Israel) and Shemi Ltd. (Shdema, Israel), and cultivar Desio was received from Mizpor Ltd. (Tquma, Israel). Cultivars White Sim, Desio and Visa belong to the standard category while cv. Eilat, Darling and Lior belong to the spray category. Stem cuttings with six or eight fully mature leaves (not counting the apical leaves which were not fully
15 expanded), harvested from greenhouse-grown plants and stored for up to 1 month at 4°C, were used to prepare stem explants.

2. Media composition and tissue-culture conditions

 Murashige and Skoog (Murashige T, Skoog F: *A revised medium for rapid growth and bioassays with tobacco tissue culture*. (1962) *Physiol Plant* 15:
20 473-497) basal medium (MS) with sucrose (30 g/l) and solidified with agar (8 g/l) (basic medium), was supplemented with growth regulators and antibiotics for cocultivation with *Agrobacterium*, regeneration and selection of adventitious shoots, and elongation and rooting of transgenic plants. All media were adjusted to pH 5.8 prior to autoclaving (121°C for 20 min). For cocultivation with
25 *Agrobacterium*, the basic medium was supplemented with 5 mg/l α -naphthalene acetic acid (NAA) and 100 μ M acetosyringone (cocultivation medium). For shoot regeneration and two-step selection of transformants, the basic medium was supplemented with 0.1 mg/l NAA and 1 mg/l 1-phenyl-3(1,2,3-thiadiazol-5-yl)-urea (TDZ) (SI-T1, first selection cycle), or with

- 11 -

Figs. 14a, 14b, 14c, 14d and 14e show flower color modification of anti-*fht* transgenic carnation. Cv. Eilat was transformed with *fht* cDNA cloned into a binary plasmid in antisense orientation 3' to 35S CaMV. a. Flower from control, non-transformed cv. Eilat. b-d. Flowers from cv. Eilat transformed with antisense *fht* (transgenic lines FHT-33, FHT-11, FHT-14, respectively). e. Complementation of FHT suppression in the petals of line FHT-11, by feeding with dihydroquercetin in 5% methanol (right). On the left, petals incubated in 5% methanol.

Fig. 15 illustrates molecular and biochemical characterization of anti-*fht* transgenic plants. a. Northern blot analysis of *fht* transcript levels in petals of anti-*fht* transgenic FHT-11, FHT-14 and FHT-33 and control, cv. Eilat. For comparison, analysis of *dfr* transcript is presented. b. Analysis of *fht* sense and antisense transcript accumulation in petals of control, cv. Eilat and anti-*fht* transgenic FHT-11 and FHT-33. Strand-specific primers were used to reverse-transcribe sense (S), antisense (A) and both (S/A) transcripts, followed by PCR amplification and Southern analysis. *chs* analysis was used as a control. c. FHT enzyme activity in petal extracts of control, cv. Eilat and anti-*fht* transgenic FHT-11 plants. [¹⁴C]naringenin (NAR) was used as a substrate for the reaction, and the FHT enzyme activity product, dihydrokaempferol (DHK), was analyzed by TLC followed by autoradiography.

Fig. 16 shows gas chromatography-mass spectroscopy (GC-MS) headspace analysis of volatile compounds, emitted from detached flowers of anti-*fht* transgenic FHT-11 plants. A representative GC chromatogram is presented. 1, methylbenzoate; 2, hexanoic acid; 3, *trans*-caryophyllene; and

Fig. 17 is a schematic presentation of anthocyanin and benzoic-acid-derivative biosynthesis, originating from the general phenylpropanoid pathway. The diversion of metabolic flow in the direction of benzoic-acid-derivative production, as a result of FHT suppression, is depicted. Enzymes are indicated as follows: chalcone synthase (CHS); chalcone flavanone isomerase (CHI); flavanone 3-hydroxylase (FHT); dihydroflavonol 4-reductase (DFR). ISP, isosalipurposide.

- 14 -

The GUS-encoding gene is not expressed in *Agrobacterium* cells carrying pKIWI105 due to the lack of a bacterial ribosome-binding site, making this plasmid suitable for transient transformation studies (Janssen, *et al*). Digestion of pCGN7001 with EcoRI releases a 3.8-kb. fragment containing uidA and part of *nptII*, whereas HindIII is a unique restriction site within the T-DNA fragment (Comai, *et al*).

Bacteria from a single colony were grown at 28°C for ca. 20 hours in liquid LB medium (10 g/l bacto-tryptone, 5 g/l bacto-yeast extract, 5 g/l NaCl, 2 g/l glucose, pH 7.5) on a rotary shaker (250 rpm). The medium was supplemented with 100 µM acetosyringone, 50 mg/l rifampicin, and 25 mg/l gentamycin or 50 mg/l kanamycin for pCGN7001 or pKIWI105, respectively. Bacteria ($OD_{550} = 0.5$) was harvested by centrifugation at 10000g for 2 min; the pellet was resuspended in liquid cocultivation medium ($OD_{550} = 0.5$ or 1.0), and the suspension was used for inoculation.

4. Microprojectile bombardment

Leaves and shoot apices were removed from sterilized cuttings and the three primary nodes were wounded prior to inoculation in a Biolistics PDS 1000/He system using tungsten M-25, 1.7 µm in diameter (Bio-Rad, Richmond, CA, USA). Explants placed on MS basal medium solidified with 1% agar (30 explants per petri dish) were bombarded twice. Each bombardment consisted of 100 µl of tungsten particles in an aqueous suspension (15 mg/ml). The particles were finely dispersed with ultrasonic cleaner (Sonicor Instrument Corporation, Copiague, NY, USA) before bombardment. Particle acceleration parameters were: a bombardment pressure of 1500 psi and a distance of 9 cm from the launching plate to the tissue.

5. Optimization of transient transformation

Stem explants with or without prior wounding by microprojectile bombardment were immersed for 10 min in a bacterial (*A. tumefaciens* EHA105/pKIWI105) suspension ($OD_{550} = 1$). Inoculated stem explants were then blotted dry and cultured in an upright position on the cocultivation medium under various light regimes for a period of up to 5 days. Following cocultivation, stem

- 13 -

0.1 mg/l NAA and 1 mg/l 6-benzylaminopurine (BAP) (SI-B1, second selection cycle). Both media were also supplemented with 300 mg/l carbenicillin and, unless otherwise stated, 100 mg/l kanamycin. Elongation and rooting of transgenic shoots, following the second selection cycle, were performed on the basic medium containing 0.1 mg/l NAA, 0.1 mg/l gibberellic acid (GA), 200 mg/l carbenicillin and 100 mg/l kanamycin. All cultures were maintained in a growth room at $25 \pm 1^\circ\text{C}$ under a 16-h photoperiod using cool white light ($60 \mu\text{mol m}^{-2}\text{s}^{-1}$) unless otherwise indicated.

3. Bacterial strains

Agrobacterium tumefaciens strains AGLO (Lazo G, Stein P, Ludwig R: *A DNA transformation-competent Arabidopsis genomic library in Agrobacterium*. (1991) Bio/Tech 9: 963-967) and EHA105 (Hood E, Gelvin S, Melchers L, Hoekema A: *New Agrobacterium helper plasmids for gene transfer to plants*. (1993) Trans Res 2: 208-218.) carrying the binary plasmids pCGN7001 (Comai L, Moran P, Maslyar D: *Novel and useful properties of a chimeric plant promoter combining CaMV 35S and MAS elements*. (1990) Plant Mol Biol 15: 373-381) and pKIWI105 (Janssen B, Gardner R: *Localized transient expression of GUS in leaf disks following cocultivation with Agrobacterium*. (1989) Plant Mol Biol 14: 61-72), respectively, were used for transient and stable transformation of carnation, respectively. Both plasmids carried the uidA gene coding for β -glucuronidase (GUS) (Stomp AM: *Histochemical localization of β -glucuronidase*. In: Gallagher SR (ed) GUS Protocols: Using the GUS Gene as a Reporter of Gene Expression, pp. 103-113. Academic Press, San Diego (1992)) driven by either a cauliflower mosaic virus (CaMV) 35S promoter (pKIWI105) or a mannopine synthetase (MAS) promoter (pCGN7001) (Comai L, *et al*), and the nptII gene coding for neomycin phosphotransferase II (NPTII) (Beck E, Ludwig G, Auerswald EA, Reiss R, Schaller H: *Nucleotide sequence and exact location of the neomycin phosphotransferase gene from transposon Tn5*. (1982) Gene 19: 227-336) driven by either a nopaline synthase (NOS) promoter (pKIWI105) or the CaMV 35S promoter (pCGN7001).

- 16 -

progeny. All crosses were carried out in the fall/winter and seeds were collected and sown in the summer. Seedlings were evaluated for GUS expression and were subjected to *uidA* and *nptII* PCR analysis. All transformation experiments were repeated at least five times.

5 7. *Evaluation of transformants*

a. GUS expression

A histochemical assay of GUS activity was performed according to Stomp. Tissue samples were incubated for a few hours to overnight at 37°C in a 0.1% (w/v) X-Gluc (5-bromo-4-chloro-3-indolyl β -D-glucuronic acid sodium salt, Biosynth Inc., Staad, Switzerland) solution containing 0.1 M sodium phosphate buffer (pH 7.0), 10 mM EDTA, and 0.1% (w/v) Triton X-100. When necessary, green tissues were bleached, after staining, by immersion in 50% (v/v) EtOH for a few hours, followed by several washes with 70% EtOH. It should be noted that no background GUS activity was detectable in any of the analyzed—intact or
15 wounded—tissues of control plants.

b. Polymerase chain reaction (PCR) analysis

DNA extraction, primers for *uidA*, and PCR conditions were as previously described (Tzfira T, Jensen CS, Wangxia W, Zuker A, Altman A, Vainstein A: Transgenic Populus: a step-by-step protocol for its *Agrobacterium*-mediated transformation. (1997) Plant Mol Biol Rep 15: 219-235). The primers for *nptII* amplification were 5'-GAGGCTATTCGGCTATGACT-3' and 5'-AATCTCGTGATGGCAGGTTG-3'. The predicted sizes of the amplified DNA fragments were 0.53 kb and 0.8 kb for *uidA* and *nptII*, respectively. Amplified DNA was electrophoresed on a 1.5% (w/v) agarose gel, using Tris-borate buffer
25 (1.3 M Tris, 0.7 M boric acid and 24.5 mM EDTA, pH 8.4). Gels were stained with ethidium bromide, photographed under ultraviolet light, and analyzed by Southern blotting.

c. Southern blot analysis

DNA was extracted from leaves by the CTAB (cetyltrimethylammonium bromide) procedure described previously (Tzuri G, Hillel J, Lavi U, Haberfeld A,
30

- 15 -

explants were histochemically evaluated for transient GUS expression by counting the number of GUS-expressing stem explants, as well as the number of blue spots per explant, under a stereo-microscope.

6. Transformation and regeneration of transgenic plants

5 Following bombardment-mediated wounding, stem explants were inoculated with bacterial (AGLO/pCGN7001) suspension ($OD_{550} = 0.5$). During cocultivation and all consecutive steps, explants were cultured in an upright position. After 5 days of culture on the cocultivation medium (3 days in the dark followed by 2 days in light), three primary nodes were sectioned into ca. 3-mm
10 slices and transferred to SI-T1 medium for shoot regeneration and the first selection cycle. It should be noted that apical meristem breakage was considered undesirable. Hence, to prevent the development of non-transformed axillary shoots, all identifiable shoot apices were removed from the stem explants prior to inoculation with bacteria.

15 After 10 days of culture, the explants were cleaned again, if needed, of the occasionally developing shoots, cross-sectioned into two halves, and transferred to fresh SI-T1 medium. After ca. 2 additional weeks, clusters of regenerated adventitious shoots were excised from the primary stem explants. Leaves from all of the shoots of each independent cluster were pulled off and cultured on SI-B1
20 medium for adventitious shoot regeneration and selection of transgenes (second selection cycle).

After 10 to 12 days, new adventitious shoots emerged from the leaf basal area. These shoots were transferred to elongation and then rooting media and evaluated as to their transgenic nature. Following hardening (Vainstein A, Fisher
25 M, Ziv M: *Shoot regeneration from petals as a basis for genetic variation and transformation*. (1992) *Acta Horti* 314: 39-45), transgenic plants were transferred to the greenhouse where they developed and flowered normally. Since cv. White Sim, like most commercial carnation varieties, is male-sterile and can only be out-crossed, T_0 White Sim plants were crossed with wild-type (non-transformed)
30 red (b.l. 13261), yellow (b.l. 13262) or white (b.l. 13263) male plants to generate T_1

- 18 -

different wounding methods (vortexing of stem explants in the presence of glass beads, sand or carborundum particles, or poking and scratching with a needle or scalpel) yielded neither efficient nor reliable transient transformation following inoculation (with or without vacuum infiltration) with *Agrobacterium* (12).

5 In contrast, when explants were wounded by microprojectile bombardment, efficient and highly reproducible transient transformation was obtained, based on both the percentage of GUS-expressing inoculated explants and the frequency of transformation events per explant. Following a 5-day cocultivation with EHA105/pKIWI105 under constant light, 90% of the cv. White Sim stem explants
10 expressed GUS when they had been wounded by bombardment, as compared to only 20% GUS-expressing explants when wounding had been effected by poking or scratching. The transient frequency events (the number of blue spots per explant) increased 7.5-fold when bombardment-mediated wounding of explants was employed (Fig. 1). It should be noted that cocultivation with *Agrobacterium* for less
15 than 5 days yielded lower transformation efficiencies (data not shown).

Different light conditions during cocultivation also strongly affected the efficiency of transient transformation. An almost threefold increase in the number of transformation events per explant was obtained when non-bombarded explants were cocultivated in the dark instead of under constant light (Fig. 1). When
20 bombardment-mediated wounding was combined with etiolation during cocultivation, the frequency of transformation events per explant increased to over 10-fold that of explants wounded by other means and cocultivated in the light (Fig. 1). However, the relatively long etiolation period led to decreased regeneration from stem explants in both *Agrobacterium*-inoculated and noninoculated (control)
25 stem explants.

To restore these explants' high regeneration potential while preserving the positive effect of etiolation on transformation, the effect of different dark/light regimes on transient GUS transformation and regeneration efficiencies was assessed. When stem explants were cocultivated with *Agrobacterium* for 3 days in
30 the dark then 2 days in the light, the high regeneration efficiency of the explants

- 17 -

Vainstein A: *DNA fingerprints of ornamental plants*. (1991) Plant Sci 76: 91-97). DNA (10 µg) was digested with HindIII or EcoRI and electrophoresed in 1% (w/v) agarose gels. DNA was transferred to a nylon membrane (Hybond N+, Amersham) by capillary blotting as previously described (Maniatis T, Fritsch EF, Sambrook J: *Molecular Cloning. A Laboratory Manual*. (1982) Cold Spring Harbor Laboratory Press, Cold Spring Harbor). PstI fragments of 2.6 kb and 1.9 kb from the binary plasmid p35SGUSINT Vancanneyt G, Schmidt R, O'Connor-Sanchez A, Willmitzer L, Rocha-Sosa M: *Construction of an intron-containing marker gene: splicing of the intron in transgenic plants and its use in monitoring early events in Agrobacterium-mediated plant transformation*. (1990) Mol Gen Genet 220: 245-250) served as probes for *uidA* and *ntpII*, respectively. The probes were ³²P-labeled by random priming (Feinberg AP, Vogelstein B: *A technique for radiolabelling DNA restriction endonuclease fragments to high specific activity*. (1984) Anal Biochem 137: 266-267), denatured and added to the hybridization mixture.

Pre-hybridization and hybridization were performed as previously described (Ben-Meir H, Vainstein A: *Assessment of genetic relatedness in roses by DNA fingerprint analysis*. (1994) Sci Hortic 58: 115-121) at 65°C for 3 h and 18 h, respectively. Post-hybridization washes consisted of two high-stringency washes in 0.45 M NaCl, 0.045 mM sodium citrate, 0.1% (w/v) SDS, 65°C, for 20 min each, followed by one wash in 0.15 M NaCl, 0.015 mM sodium citrate, 0.1% SDS, 65°C, for 20 min. The blots were exposed to an imaging plate (Fujix Bas 1000, Fuji, Japan) for 2-7 h. The plate was then read in an imaging plate reader (Fujix Bio Imaging Analyzer Bas 1000). Two washes of 0.6 M NaCl, 0.06 mM sodium citrate, 0.1% SDS, 65°C, for 20 min each, were added prior to the three post-hybridization washes when Southern blots were performed with the PCR products.

B. Results

1. Optimization of transient transformation

uidA (GUS) reporter gene expression was used to monitor early transformation events in carnation stem explants. Preliminary experiments testing

- 20 -

throughout all analyzed tissues. Since the application of a higher kanamycin concentration (120 and 150 mg/l) in the first selection cycle led to reduced shoot regeneration while transformed plants remained mostly chimeric in nature (data not shown), a second selection cycle was performed to eliminate the chimeric plants.

Leaves originating from individual clusters were cultured, separately for each cluster, on SI-B1 medium to eliminate the possibility of generating transgenes representing a single transformation event. After ca. 2 weeks of the second-selection cycle, two-thirds of the independent clusters selected on 100 mg/l kanamycin yielded scorable shoots (Table 1). These adventitious shoots regenerated directly from the basal part of the leaves, and only from an area which remained green under kanamycin selection (Fig. 3D).

– 19 –

was restored, reaching the level of those cocultivated for 5 days in the light. Moreover, transformation efficiency was not affected relative to explants cocultivated for 5 days in the dark (cv. White Sim in Figs. 1 and 2).

To assess the suitability of the combined wounding and dark/light regime to other carnation cultivars, transient transformation frequencies in five varieties representing both standard and spray categories were analyzed. All five were susceptible to transformation with EHA105/pKIWI105 under the aforementioned conditions. For all cultivars, 80-90% of the inoculated explants expressed GUS, albeit with some variation in the frequency of transformation events per stem explant (Fig. 2). Further experiments were performed using this combination of wounding and inoculation/cocultivation conditions.

2. Stable transformation and regeneration of transgenic plants

To improve the effectiveness of kanamycin for transgene selection, stable transformation of cv. White Sim was performed with pCGN7001 which carries 35S-driven *nptII*, rather than the NOS-driven *nptII* of pKIWI105 used in the transient transformation experiments. When inoculation was performed with a high concentration of bacteria ($OD_{550} \geq 1$), their extensive growth prevented the establishment of aseptic cultures, even at high concentrations of antibiotics, and negatively affected adventitious shoot regeneration. Inoculation of explants with bacteria at an OD_{550} of 0.5 was optimal, allowing controlled bacterial growth with no adverse effect on the further tissue culture and regeneration of plantlets following transfer to the regeneration/selection SI-T1 medium.

After ca. 1 month in culture following inoculation, adventitious shoot clusters, regenerated directly from sectioned stem explants, were easily scorable (Fig. 3). Interestingly, while all three internodes showed high regeneration ability in aseptic tissue culture, only the two top internodes retained this potential following inoculation and selection. Histochemical evaluation of regenerated clusters revealed a chimeric GUS expression pattern in most of the adventitious shoots (Fig. 3), despite high selection pressure (100 mg/l kanamycin) during regeneration; only 1 to 3% of all regenerated shoots expressed GUS in and

- 22 -

In almost all of these shoots, histochemical assay revealed GUS expression in all organs and throughout the tissues, with no observable chimerism (Table 1, Fig. 3E, 3F). To assess the overall efficiency of the two cycles of selection, only one GUS-expressing shoot per individual cluster was counted, even though 5-20 GUS-expressing shoots were usually generated from leaves of each cluster. Based on this consideration, which allows an estimation of independent transformation events, the overall yield of the procedure was 19 GUS-expressing shoots generated per 100 *Agrobacterium*-inoculated stem explants.

The possibility of further increasing the overall yield by lowering the selection pressure during the first selection cycle was assessed (Table 1). Following cocultivation, stem explants were cultured on SI-T1 medium containing 80 and 90 mg/l kanamycin instead of 100 mg/l. The lowest kanamycin concentration yielded ca. three times more and 90 mg/l kanamycin ca. two times more shoot clusters relative to the number regenerated on 100 mg/l kanamycin. However, only ca. one-third of the clusters generated at both 80 and 90 mg/l kanamycin yielded adventitious shoots from leaves following the second selection cycle. Thus, while the overall number of independent GUS-expressing shoots was essentially the same for the three levels of kanamycin analyzed, 100 mg/l was considered optimal because it almost completely prevented the generation of escapees (Table 1).

Forty randomly selected independent, kanamycin-resistant, GUS-expressing T₀ cv. White Sim plants exhibited a normal phenotype when, following hardening, they were grown to flowering in the greenhouse. Progeny generated from an outcross of the T₀ White Sim lines also expressed GUS and flowered normally (Fig. 3G, 3H).

To further confirm the transgenic nature of the generated kanamycin-resistant GUS-expressing plants, Southern blot analysis was performed. Hybridization of *Eco*RI-digested genomic DNA from T₀ lines with *uidA* probe yielded the expected 3.8 kb fragment (Fig. 4, lanes 1-4); this fragment was not detectable in the non-transformed control line (Fig. 4, lane C). Two to

- 21 -

Table 1.

Efficiency of selection of transformed carnation plants. Stem explants of cv. White Sim. were transformed with *Agrobacterium tumefaciens* AGLO/pCGN7001 as described in the experimental protocol.

First selection cycle		Second selection cycle (100 mg/l kanamycin)	
Kanamycin (mg/l)	No. of selected clusters *	No. of clusters generating shoots **	No. of GUS- expressing shoots ***
80	92±35	29±1 (32)	17±1 (59)
90	65±4	25±2 (38)	20±1 (80)
100	32±4	21±1 (66)	19±1 (90)

* The number of harvested independent shoot clusters regenerated from a total of 100 stem explants, following the first selection cycle.

10 ** The number of independent clusters that generated shoots from leaves following the second selection cycle. The numbers in brackets represent percentages of independent clusters generating shoots from leaves following the second selection cycle out of the total number of independent clusters selected in the first selection cycle.

15 *** The number of independent clusters generating GUS-positive shoots from leaves following the second selection cycle. The numbers in brackets represent percentages of independent clusters generating GUS-positive shoots from leaves out of the total number of independent clusters generating shoots following the second selection cycle.

- 24 -

K.R. Summerfelt. 1990. Improved binary vectors for *Agrobacterium*-mediated plant transformation. Plant Mol. Biol. 14:269-276.) binary plasmid to create the binary plasmid pAMrolC. pAMrolC contains the *nptII* selection gene (coding for neomycin phosphotransferase II (*NPTII*); Beck, E., G. Ludwig, E.A. Auerwald, R. Reiss, and H. Schaller. 1982. Nucleotide sequence and exact location of the neomycin phosphotransferase gene from transposon Tn5. Gene 19:227-336) also driven by a 35S promoter. The plasmids pCGN7001 (Comai et al., 1990) (a control plasmid essentially identical to pCGN1559, but carrying the *uidA* gene coding for β -glucuronidase (GUS) driven by a mannopine synthetase (MAS) promoter and the *nptII* gene driven by the 35S promoter) and pAMrolC were transferred to *A. tumefaciens* strain AGLO (Lazo, G., P. Stein, and R. Ludwig. 1991. A DNA transformation-competent *Arabidopsis* genomic library in *Agrobacterium*. Bio/Technology 9:963-967) and used to transform *Dianthus caryophyllus* L. cv. White Sim as described above. Eighteen independent pAMrolC transformants were obtained, of which three *rolC*-transgenic lines (R-1, R-2 and R-3) were studied in more detail. The transgenic origin of the *uidA*-expressing control lines, which were phenotypically and morphologically indistinguishable from non-transformed plants, was previously confirmed as described above.

2. Southern and northern blot analyses.

The cetyltrimethylammonium bromide (CTAB) extraction procedure and a TRI-REAGENT™ extraction kit (Molecular Research Center, Inc., Cincinnati, OH) were employed for DNA and RNA extraction, respectively, from leaves of greenhouse-grown plants, as described previously. DNA (10 μ g) was digested with *HindIII* and electrophoresed through a 1% agarose gel using Tris-borate buffer (containing 1.3 M Tris, 0.7 M boric acid and 24.5 mM EDTA, pH 8.4). *HindIII* is a unique restriction site within the T-DNA fragment, 5' to *rolC*. Gels were stained with ethidium bromide and photographed under ultraviolet light. Total RNA (10 μ g) was electrophoresed through a 1.2% formaldehyde gel (Vishnevetsky, M., M. Ovadis, H. Izhaki, M. Levy, Y. Libal-Weksler, Z. Adam, and A. Vainstein. 1996. Molecular cloning of a carotenoid-associated protein from *Cucumis sativus*

- 23 -

four fragments of different sizes were revealed following digestion of T₀ genomic DNA with *Hind*III (Figure 4), thus confirming integration of the GUS-encoding gene construct in the plant genome. Integration of the *nptII* gene into the T₀ plant genome was also confirmed by Southern blotting using *nptII* as a probe (data not shown). The molecular analysis of seedlings derived from one of the crosses between T₀ plants and non-transgenic breeding line (b.l.) 13261 is shown in Fig. 5. *nptII* PCR amplification followed by Southern blotting using *nptII* as a probe yielded a DNA fragment of the expected size (0.8 kb) in all analyzed kanamycin-resistant T₁ seedlings (Fig. 5) and not in controls (Fig. 5, lane C). Similarly, *uidA* PCR amplification of these T₁ seedlings followed by hybridization with *uidA* probe yielded a DNA fragment of the expected 0.53 kb in all cases (data not shown).

The applicability of the transformation procedure was also assessed with another two, genetically unrelated, commercially highly successful cultivars—Desio and Eilat, which had been used in transient transformation experiments. The efficiency of selection for these varieties was essentially identical to that detailed for cv. White Sim and overall transformation efficiencies of 13% and 18% (transgenes out of total inoculated explants) were obtained for cvs. Desio and Eilat, respectively. The transgenic nature of these plants was confirmed by Southern blot analysis (data not shown). Interestingly, the overall efficiencies of stable transformation for the three analyzed varieties were in good agreement with the frequency of transient transformation events (see Fig. 2).

II. *rolC* TRANSGENIC CARNATION

A. Materials and Methods

1. Gene construct and plant genetic transformation.

A 1.46-kb *Xba*I fragment from plasmid pUC18-CaMV35S-*rolC* (13), carrying the *A. rhizogenes rolC* gene (Fig. 6) driven by the CaMV 35S RNA promoter, was inserted into the *Xba*I-predigested pCGN1559 (McBride, K.E. and

- 26 -

Stems cuttings dipped in Hormoril® and rooted in peat mixture were also transferred to no. 0.8 volcanic rock mixture for growth under regular greenhouse conditions, in order to establish mother plants for further clonal propagation. Four weeks after transfer to the greenhouse, plants were decapitated to allow for axillary bud development. Stem cuttings with at least three internodes, originating from the breaking and elongation of the axillary buds, were then harvested from the mother plants (40 plants per line) ca. every 4 weeks. A total of five harvest cycles was performed and the number of stem cuttings determined. In addition, the number of internodes in the stem cuttings obtained from the first harvest cycle was determined. Side shoots developing on these cuttings were also characterized as to their dry weight and the number of open leaves (Fig. 7).

Plant height at flowering was measured from the soil line to the top of the main flower bud on the flowering stem. The number of petals per flower bud, their length and dry weight were measured from flowers at anthesis, during the flowering season (Nov 98 – Mar 99). The vase life of flowers with a 50-cm-long stem was measured in distilled water, at 20°C, under a 12-h photoperiod using cool white light ($60 \mu\text{mol m}^{-2}\text{s}^{-1}$).

B. Results

1. Confirming the transgenic nature of selected lines.

Transgenic plants were generated following two regeneration and selection cycles under high kanamycin pressure as described previously. All *rolC*- and control, *uidA*-transgenic plants exhibited high resistance to kanamycin, and the latter also exhibited strong GUS expression in all their organs. *rolC*-transgenic plants (18 independent lines) were screened *in vitro* for phenotypic alterations which included autonomous root growth in hormone-free media, axillary bud breakage, increased growth rate and altered internode length. Out of 18 lines exhibiting altered morphology, three independent lines (R-1, R-2 and R-3) were selected, hardened and transferred to the greenhouse for further study. Southern blot analysis of *HindIII*-digested DNA revealed integration of the *rolC* gene in all three selected lines (Fig. 8A). Northern blot analysis revealed the expression of

- 25 -

corollas: homologous genes involved in carotenoid sequestration in chromoplasts. Plant J. 10:1111-1118.). DNA and RNA were transferred to a nylon membrane (Hybond N⁺; Amersham, UK) by capillary blotting as described by Maniatis et al. (1982). The 0.76-kb *Sa*II fragment from plasmid pUC18-CaMV35S-*rolC*,
5 representing the 3' prime end of the *rolC* gene, served as a probe for that gene. Probe-labeling, and prehybridization and hybridization of Southern and northern blots were performed as described above. The blots were visualized by exposure to Agfa Curix PR2 film at -70°C for 5 to 20 h with an intensifying screen.

3. *Hardening of tissue-culture plants and establishment of plants in the*
10 *greenhouse.*

About 3-cm long, rooted plantlets were removed from the *in vitro* culture medium and their roots were cleaned of agar. The plantlets were then transferred to plastic planters filled with water-saturated no. 4 perlite (Agrical, Kibbutz Habonim, Israel) and kept for 2 weeks on a heated rooting table in a greenhouse under
15 intermittent mist (misting cycles of approximately 3 min every half hour). The plantlets were then transferred to a no. 0.8 volcanic rock (Tof Marom Golan, Marom Golan, Israel) mixture and further grown for 4 weeks in the same rooting greenhouse. Hardened plants were then transferred to a regular greenhouse and 4 weeks later they were decapitated to allow breaking and elongation of axillary
20 shoots. These plants served as the source for standard stem cuttings for all further experiments.

4. *Assessment of developmental traits in the greenhouse.*

To evaluate the rooting performance of transgenic plants, stem cuttings, either directly or following dipping in 0.6% (w/w) indole-3-butyric acid (IBA)
25 containing commercial rooting powder (Hormoril® T-6, Asia Rizel, Ramat Gan, Israel), were planted in either no. 4 perlite or a rooting mixture (50% v/v no. 4 perlite, 10% v/v no. 2 perlite, 20% v/v slurry and 20% v/v peat). The stem cuttings (50 cuttings per line) were grown for 3 weeks on heated rooting tables. The cuttings planted in perlite only were then gently pulled and washed of rooting
30 mixture. The root system was dissected, dried at 65°C for 48 h and weighed.

- 28 -

To further characterize type IV and V stem cuttings, they were dissected and side shoots, developed from axillary buds, were weighed and their developmental stage ranked (Fig. 7, Table 2). Side shoots of *rolC*-transgenic stem cuttings exhibited an up to 1.5-fold higher dry weight than those originating from control plants (Table 2). Relative to control plants, the side shoots from *rolC* transgenes were much more developed (Table 2). In *rolC* transgenes, the average number of open leaf pairs on the side shoots at the 5th position ranged from 1.4 to 2.0. In line R-1, all side shoots at the 5th position had two open pairs of leaves. Furthermore, side shoots at the 4th position in all *rolC* transgenes averaged at least one pair of open leaves. In contrast, in control plants all side shoots developing at the 4th or 5th positions were without open leaves (Table 2).

3. Rooting ability of *rolC*-transgenic carnation plants.

When stem cuttings were not treated with auxin (Hormoril® rooting powder), control (GUS-transgenic) cuttings yielded short and undeveloped adventitious roots, whereas *rolC*-transgenic cuttings developed a dense root system with a high dry weight mass, ca. two to five times higher than that of the control cuttings (Figs. 10, 11). Moreover, the percentage of cuttings developing roots out of the total number of cuttings was ca. two times higher in R-1 and R-3 plants than in controls. In fact, nontreated stem cuttings of line R-1 exhibited root-system structures and dry weights similar to those of control plants treated with rooting powder (Figs. 10, 11). The enhanced rooting ability of *rolC* transgenes vs. controls was also apparent following treatment of the cuttings with Hormoril®: *rolC* transgenes developed up to twice the root mass of control cuttings. Moreover, while a period of ca. 3 weeks was essential for the rooting of control carnation stem cuttings, a shorter period of only 2 weeks was required before *rolC*-transgenic carnation stem cuttings could be transferred for further development to standard greenhouse conditions.

4. Plant growth and flower development.

To further assess the growth habits of *rolC*-transgenic and control (GUS-transgenic) plants, they were grown to flowering in the greenhouse. *rolC*

- 27 -

rolC, driven by the constitutive 35S promoter, in these three lines (Fig. 8B), thus confirming their transgenic nature.

2. Yield and morphology of stem cuttings from *rolC*-transgenic vs. control plants.

Stem cuttings were collected from *rolC*- and GUS-transgenic (control) carnation plants and classified, by developmental stage, based on the number of internodes per cutting (Fig. 7). About 50% of all stem cuttings from GUS-transgenic plants were of type III (three internodes), and only ca. 6% were of type V. In contrast, up to 97% of the *rolC*-transgene cuttings were of types IV and V (Fig. 9). Furthermore, the overall yield of cuttings per mother plant was up to 40% higher in *rolC* transgenes than in control plants (Table 2).

Table 2

Stem-cutting yield and properties in *rolC*-transgenic and control carnation plants.

Line	Stem cuttings	Side shoots		
	Yield ^a	Dry weight ^b	Open pairs of leaves ^c	
	(no.)	(positions 4 and 5) (mg)	position 4	position 5
R-1	23 ± 2	47 ± 1	1.6 ± 0.1	2.0 ± 0
R-2	21 ± 1	44 ± 2	1.6 ± 0	1.8 ± 0.1
R-3	21 ± 1	36 ± 1	1.1 ± 0.1	1.4 ± 0.2
C	16 ± 2	29 ± 3	0	0

^a Stem cuttings were harvested a total of five times at 3- to 5-week intervals. An average number of stem cuttings ± SE per harvest per mother plant is presented.

^b Average weight of side shoot ± SE, at positions 4 and 5, per stem cutting.

^c The average number of open pairs of leaves ± SE per side shoot at positions 4 or 5 is presented.

- 30 -

fragment was subcloned in antisense orientation (Fig. 13) between the CaMV 35S promoter and *nos* terminator in pJD330 (Broido, S., Loyter, A. & Vainstein, A. Transient expression of photosynthetic genes in transfected albinoid petunia protoplasts and correct processing of newly synthesized chloroplast-destined polypeptides. *Physiol. Plant.* 88, 259-266 (1993)) to create pJD-anti-*fht*. An *Xba*I fragment from pJD-anti-*fht* was then cloned into the binary vector pCGN1559 (Comai, 1990) to create pAM-anti-*fht*. The gene construct was transferred to carnation cv. Eilat via cocultivation of stem explants, as described above, with *Agrobacterium tumefaciens*, AGLO containing pAM-anti-*fht*.

10 2. Molecular and biochemical analysis.

For northern analysis, RNA was extracted from petals and samples (10 µg) were electrophoresed through an agarose gel, transferred to a membrane and probed with ³²P-labeled *fht* and *dfr* cDNAs as described previously (Vishnevesky, M. et al). For RT-PCR analysis (de Lang, P., de Boer, G.-J., Mol, J.N.M. & Kooter, M. Conditional inhibition of β -glucuronidase expression by antisense gene fragments in petunia protoplast. *Plant Mol. Biol.* 23, 45-55 (1993)), RNA samples treated with DNase were reverse-transcribed with M-MLV reverse transcriptase, using strand-specific *fht* and *chs* (sense and antisense) primers. The resultant cDNAs were PCR-amplified using a mixture of both sense and antisense primers. The amplified product was analyzed by Southern blotting as described above, using ³²P-labeled *fht* and *chs*.

Carnation cDNA clones of *chs* and *dfr* were isolated, as described above for FHT, by PCR using specific primers according to their sequences in GenBank: *chs* (Z67982), *dfr* (Z67983). Primers used were: 5' CCC AAA ACG CTC ACT TCA CT 3' and 5' CCA AGC CCA TCT AAG CAA GT 3' for *fht*; 5' GGG CCG ATG GTC CTG CTA CTA T 3' and 5' ACG CGC TCG ACA TGT TCC CAA A 3' for *chs*; 5' TGT GAA TGT CGA AGC GAC TC 3' and 5' TTG AAT TTG GTG GGG ACA TT 3' for *dfr*.

Feeding experiments (Deroles, S.C. et al. An antisense chalcone synthase cDNA leads to novel colour patterns in lisianthus (*Eustoma grandiflorum*) flowers.

- 29 -

transgenes developed a higher number of shoots and a more elaborate root system than control plants (Fig. 11). Furthermore, during the 5-month flowering season, the average number of flowering stems per mother plant in R-3 *rolC* transgenes was three times that in control plants (9.4 ± 1.8 , 3.2 ± 1.0), two and two-and-a-half times for R-1 and R-2 (7.2 ± 1.4 , 8.1 ± 1.7). The average number of flower buds on flowering stems was essentially the same (4.5 ± 0.5 flowers/stem) in *rolC* and control plants.

The height of the flowering stems of *rolC* transgenes was similar to that of control plants, being 85 ± 5 , 87 ± 4 and 95 ± 6 vs. 75 ± 5 cm in R-1, R-2, R-3 and control plants, respectively. Time to flowering was essentially identical in *rolC* transgenes and control plants, ca. 14 weeks after decapitation of mother plants. Flower vase life of control and *rolC* transgenes was also the same (12 ± 2 days). An assessment of flower size during the flowering season revealed that R-1 and R-3 transgenic flowers are essentially identical to control ones, in all analyzed parameters, i.e. dry weight, petal number and size. R-2 transgene flowers were smaller (Fig. 11) due to a smaller number of petals: 410 ± 54 mg flower dry weight with 35 ± 3 petals (24 ± 3 mm in length) as compared to 696 ± 53 mg dry weight with 55 ± 2 petals (33 ± 3 mm in length) in control flowers. Crosses between R-1, R-2, R-3 and non-transformed male carnation plants revealed that introduction of the *rolC* gene does not affect the flower's female fertility (not shown). The effect of *rolC* on male fertility could not be tested, because White Sim, the cultivar used in this study to generate transgenes, is male-sterile.

III. ANTISENSE SUPPRESSION OF *fht*

A. Materials and Methods

25 1. Plant transformation and vector construction

The cDNA sequence of the *fht* gene is shown in Fig. 12. A carnation partial *fht* cDNA clone (1 kb) was isolated by PCR of reverse-transcribed poly (A)⁺ RNA, using specific primers according to the sequence in GenBank (X70378). This

- 32 -

Evaluation Techniques (CRC Press Inc, Boca Raton, FL, 1991)) (1-5 scale, 1 for no fragrance, 5 for very intense fragrance) was performed with 20 panelists.

B. Results

The flowers of carnation cv. Eilat have dark orange petals with reddish edges (Fig. 14a). The enzyme flavonoid 3',5'-hydroxylase is lacking in carnation, and in cv. Eilat, flavonoid 3'-hydroxylase (F3'H) is inactive: only pelargonidin and the yellow isosalipurposide (ISP) are accumulated. Following transformation of cv. Eilat with antisense *fht*, 14 individual transgenic plants were regenerated and grown to flowering in the greenhouse. All plants developed and flowered normally and six of them exhibited color modifications ranging from attenuation to complete loss of the orange/reddish color (Fig. 14b-d). A complete loss of orange/reddish color and the production of yellow/cream flowers, accumulating ISP and only traces of pelargonidin, was observed in two independent anti-*fht* transformants (FHT-11 and FHT-14, Fig. 14c and 14d, respectively). Transformation of cv. Eilat with the identical vector carrying the *gus* reporter gene instead of *fht* led only to transgenes with true-to-type flower color.

Since, like many commercial carnation varieties, cv. Eilat is sterile, the inheritance of color modification could not be analyzed. Southern blot analysis confirmed the presence and integration of anti-*fht* gene construct into the genome of the transgenic plants (data not shown). Northern blot analysis (Fig. 15a) revealed dramatic suppression of *fht* transcript in transgenes FHT-11 and FHT-14, in contrast to controls and transgene FHT-33. To further detail the suppression of the *fht* gene, RT-PCR analysis of the endogenous and transgenic *fht* transcripts was performed (de Lang, et al, 1993) using strand-specific primers. Petals of FHT-11 transgenes, which exhibited a dramatically modified phenotype (Fig. 14c), did not accumulate detectable levels of sense *fht* transcript (Fig. 15b). In contrast, petals of transgenic line FHT-33, which showed only minor color alterations (Fig. 14b), accumulated relatively high levels of sense transcript. The level of *fht* antisense transcript was much lower in FHT-11 than in the FHT-33 transgene. In control flowers, as expected, only sense transcript could be detected. Analysis of *chs*

- 31 -

Mol. Breed. 4, 59-66 (1998)) were performed with detached petals, incubated for several hours in 1 mg/ml dihydroquercetin in 5% methanol. FHT activity was assayed using petal extracts (Britsch, L. & Grisebach, H. Purification and characterization of (2s)-flavanone 3-hydroxylase from *Petunia hybrida*. *Eur. J. Biochem.* 156, 569-577 (1986)) and flavonoid analysis was performed by TLC (Harborne, L.B. *Comparative Biochemistry of the Flavonoids* (Academic Press, New York, 1967)).

3. Headspace GC-MS analysis and fragrance tests.

Detached flowers at anthesis were sealed in a 35-ml glass tube for 30 min and volatile compounds were trapped for 10 min using polydimethylsiloxane-coating solid-phase-microextraction (SPME) fiber assembly (Supelco Inc., Bellefonte, PA). As a blank, trapping was performed in sealed tubes without flowers. Compounds were analyzed by SPME headspace GC-MS (Helsper, J.P.F.G., Davies, J.A., Bouwmeester, H.J., Krol, A.F. & van Kampen, M.H. Circadian rhythmicity in emission of volatile compounds by flowers of *Rosa hybrida* L. cv. Honesty. *Planta* 207, 88-95 (1998)) (Varian Star 3400 CX GC equipped with a 30 m x 0.25 mm ID, DB-5MS column and interfaced with a Varian Saturn 3 MS). The fiber was manually placed into a 1077 splitless injector for 3 min (injector temperature was 210°C). Helium served as the carrier gas and GC temperature was programmed as follows: 3 min at 40°C, increased to 220°C at 15°C min⁻¹ and then 5 min at 220°C. Ionization energy was 70 eV. Each compound was tentatively identified (>95% match) based on the Wiley library (Wiley registry of mass spectral data, 6th edition, F. W. McLafferty, 1994, J. Wiley & Sons, Inc). To confirm identification, standard compounds were used to match GC retention times and MS spectra. Authentic compounds were obtained from Sigma Chemical Co. (St. Louis, MO).

Paired-comparison fragrance test (O'Mahony, M. *Sensory Evaluation of Food* (Marcel Dekker, New York, 1986)) was performed with 45 panelists. An intensity category scaling test (Meilgaard, M., Civille, G.V. & Carr, T.B. *Sensory*

- 34 -

environmental conditions and sampling times) in flowers of FHT-11 and 14 relative to controls. The levels of *trans*-caryophyllene and hexanoic acid were not affected by the anti-*fht* transgene. Hence, essentially complete inhibition of dihydroflavonol biosynthesis by *fht* antisense diverts metabolic flow towards the biosynthesis of benzoic acid derivatives, which also originate from the phenylpropanoid pathway (Fig. 17). Methylbenzoate is a known fragrance compound produced by many flowers, and therefore it is not unreasonable to propose that this compound is responsible, at least in part, for the enhanced fragrance in the transgenic plants. To this end, both flower fragrance and methylbenzoate levels were increased in control plants following feeding with the *fht* substrate, naringenin (not shown).

- 33 -

transcript, used as a control, yielded the expected results, i.e. similar levels of sense transcript were detected in all analyzed plants. Analysis of FHT enzyme activity in petals of FHT-11 revealed no detectable activity, in contrast to strong FHT activity in control plants (Fig. 15c). This blockage in the anthocyanin biosynthetic pathway could be complemented by feeding detached petals with the product of the FHT enzyme, dihydroquercetin (Fig. 14e).

During the course of a 2-year field test, we observed that flowers of two independent transgenic plants (FHT-11 and 14) were more fragrant than non-transformed or GUS-transgenic controls. The enhanced fragrance was observed in flowers with strongly reduced anthocyanin accumulation, but not in FHT-33 or other transgenic lines exhibiting less dramatic reductions in anthocyanin levels. To confirm the observation that the anti-*fht* transgenes (FHT-11 and FHT-14) were more fragrant than controls, a paired-comparison fragrance test was performed. The overall fragrance intensity of transgenic FHT-11 flowers was significantly higher ($p = 0.001$) than that of control flowers. Furthermore, an intensity category scaling test revealed a highly significant difference ($p < 0.001$, paired t-test) in the intensity scores of transgenic FHT-11 (mean \pm SE = 3.2 ± 0.3) vs. control (1.3 ± 0.2) flowers. Essentially identical results were found for transgene FHT-14.

To further characterize the scent of transgenic and control flowers, gas chromatography-mass spectroscopy (GC-MS) headspace analyses were performed. Cv. Eilat and transgenic plants FHT-11 and 14 were grown in the greenhouse side by side, and flowers at anthesis were collected simultaneously and immediately analyzed. Among the major volatiles generated by carnation flowers, various known fragrance compounds were found. These compounds represented different metabolic pathways: the terpenoid *trans*-caryophyllene, the fatty acid derivative hexanoic acid, and the benzoic acid derivative methylbenzoate (Fig. 16).

In all analyses performed during the 4-month-flowering period (20 independent analyses, performed on different dates, with four flowers per line), the level of methylbenzoate was higher by a factor of 10 to 100 (varying with

- 36 -

10. A transgenic carnation plant according to Claim 9 wherein said DNA molecule is the *Agrobacterium rhizogenes rolC* gene.
11. A plant according to Claim 10 wherein said *rolC* gene is under the control of a cauliflower mosaic virus (CaMV) 35S promoter.
- 5 12. A transgenic carnation plant transformed with the *Agrobacterium rhizogenes rolC* gene.
13. A transgenic carnation plant transformed with a DNA molecule comprising the sequence of Fig. 6.
14. A transgenic carnation plant according to Claim 9 wherein expression of
10 the flavanone 3-hydroxylase (*fht*) gene is blocked.
15. A plant according to Claim 14 wherein the expression of the *fht* gene is blocked by an antisense molecule.
16. A plant according to Claim 15 wherein said antisense molecule comprises the whole or a fragment of the *fht* gene in antisense orientation.
- 15 17. A plant according to Claim 16 wherein said antisense molecule comprises the sequence of Fig. 13.
18. A transgenic carnation plant wherein expression of the flavanone 3-hydroxylase (*fht*) gene is blocked.
19. A transgenic carnation plant transformed with a DNA molecule comprising
20 the sequence of Fig. 13.
20. A DNA sequence of Fig. 13.
21. A method for controlling the fragrance of a plant comprising modulating gene expression in the anthocyanin-biosynthetic pathway of said plant.
22. A method according to Claim 21 where said plant is a carnation.
- 25 23. A method according to Claim 21 wherein said modulation comprises suppressing the expression of one or more genes in said pathway.
24. A method according to Claim 23 wherein said modulation comprises suppressing the expression of the *fht* gene.
25. A method according to Claim 24 wherein expression of the *fht* gene is
30 suppressed by an antisense molecule.

- 35 -

CLAIMS:

1. A method of transforming a carnation (*Dianthus L.*) plant genome with a DNA molecule comprising:
 - (a) preparing stem explants from carnation cuttings;
 - 5 (b) wounding the explants by microprojectile bombardment;
 - (c) cocultivating said wounded explants with *Agrobacterium* comprising said DNA molecule under conditions of exposure to dark followed by light;
 - (d) excising shoots from said cultivated wounded explants and removing
10 the leaves from said shoots; and
 - (e) culturing said leaves to obtain transgenic shoots transformed with said DNA molecule.
2. A method according to Claim 1 wherein said conditions of dark followed by light comprise 2-4 days in the dark followed by 1-4 days in the light.
- 15 3. A method according to Claim 1 wherein said carnation cuttings are stem cuttings with four to ten fully mature leaves.
4. A method according to Claim 3 wherein leaves and shoot apices are removed from said stem cuttings and two to five primary nodes are isolated.
5. A method according to Claim 1 wherein said microprojectile bombardment
20 comprises accelerating tungsten particles at said stem explants.
6. A method according to Claim 5 wherein said tungsten particles are accelerated at a pressure of 1300-2000 psi and at a distance of 3-12 cm from said explants.
7. A method according to Claim 1 wherein said wounded explants are
25 cocultivated with *Agrobacterium* in a medium supplemented by α -naphthalene acetic acid (NAA) and 1-phenyl-3(1,2,3-thiadiazol-5-yl)-urea (TDZ).
8. A method according to Claim 1 wherein the leaves of said excised shoots are cultured in a medium supplemented by NAA and 6-benzylaminopurine (BAP).
9. A transgenic carnation plant obtained by the method of Claim 1.

1/18




Fig. 1

- 37 -

26. A transgenic plant wherein the fragrance of said plant has been changed by modulating gene expression in the anthocyanin-biosynthetic pathway of said plant.

27. A method for controlling the color of a plant comprising modulating gene expression in the anthocyanin-biosynthetic pathway of said plant.

5 28. A method according to Claim 27 wherein said plant is a carnation.

3/18




Fig. 3

2/18




Fig. 2

5/18




Fig. 5

800bp

4/18




Fig. 4

7/18

901 TCATAGACCA TGCAAAGTCA TACTACCTGA AACTGATAAA TAACGACAAA GAAAGTAGCC

961 TATTAAAAG TCGCTATAGC ATGAATT

Fig. 6 (cont.)

6/18

1 ATGGCTGAAG ACGACCTGTG TTCTCTCTTT TTCAAGCTCA AAGTGGAGGA TGTGACAAGC
61 ACGGATGAGC TAGCTAGACA CATGAAGAAC GCCTCAAATG AGCGTAAACC CTTGATCGAG
121 CCGGGTGAGA ATCAATCGAT GGATATTGAC GAAGAAGGAG GGTCGGTGGG CCACGGGCTG
181 CTGTACCTCT ACGTCGACTG CCCGACGATG ATGCTCTGCT TCTATGGAGG GTCCTTGCCT
241 TACAATTGGA TGCAAGGCGC ACTCCTCACC AACCTTCCCC CGTACCAGCA TGATGTGACT
301 CTCGATGAGG TCAATAGAGG GCTCAGGCAA GCATCAGGTT TTTTCGGTTA CGCGGATCCT
361 ATGCGGAGCG CCTACTTCGC TGCAATTTCT TTCCCTGGGC GTGTCAATCA GCTGAATGAG
421 CAGATGGAGC TAACTTCGAC AAAGGGAAG TGCTTGACAT TCGACCTCTA TGCCAGCACC
481 CAGCTTAGGT TCGAACCTGG TGAGTTGGTG AGGCATGGCG AGTGCAAGTT TGCAATCGGC
541 TAATGGTTAG TCGATGGGCT GACGAGTTTG ATGTCAGGAG AAGCTGAGTG TGCTACTTGT
601 TTCCCTTTAA GAAGTATTAA TGTAATAAAA ATCAAGATCT GGTTTAATAA CTGGATACTT
661 GATTTCATCG CGCTTTTTTT GAATAAATGT TTGTTGTCTT GACTTTAAGA TATCCTTTGA
721 AATTGCGTT ATTGCTATTT CGCTTTTGGT TATTCCAAA AGACTTTGCT CAGTAAGATC
781 AAACGTTTGT ATTTCTCCGG GCCACAATAT TTGACCTATA TGCACCTGGC CACGCGCCGC
841 AATAGATGAA AATTGCCAAA ATTAGCTATC GGTCTTCTGA AAAGAAGGGC CGACATGTTT

Fig. 6

9/18




Fig. 8

8/18




Fig. 7

11/18




Fig. 10

10/18




Fig. 9

13/18

1 ACTATATCTT AAATATTCAC AACATTATAA CATAAGCTTC AAAATAACAT
51 TATTCGGATA TTTACGTAAT ATAATACGTA TCATATTAGG GTACATTCAT
101 TTTATCAACT ACGACTGCAT ATTGTTAGAC AGTCTCATAT ATACGCATAA
151 AAAATGGTCG CTGAAAAACC CAAAACGCTC ACTTCACTAG AAGGGGACGA
201 TAAATTGAAC TCGAATTTTG TTAGGGACGA GGATGAACGT CCGAAAGTGG
251 CGTATAATGA GTTTAGCAAT GATATTCCGG TGATATCTCT TGCTGGTATA
301 GATGGTGAAA AAAGGGGTGA AATATGTCGG AAGATTGTTG AGGCGTGTGA
351 AGATTGGGGG ATTTTCAAG TGTTGATCA CGGTGTTGGT GACGATCTTA
401 TTGCTGATAT GACTCGGTTG GCTCGTGAAT TTTTCGCTCT CCCGGCAGAA
451 GAGAAGCTCC GATTTGATAT GTCTGGTGGT AAAAAGGGCG GTTTTATCGT
501 GTCGAGTCAT CTTCAAGTTCAATCAACAGG GAGAAGTAGT GCAGGACTGG
551 AGGGAAATCGTGACGTATTT CTCATACCCG ACGAACTCAA GGGACTACAC
601 AAGATGGCCAGACAAACCAG AGGGTTGGAT AAAGGTCACA GAGGAATACA
651 GCAACAAGTTAATGACCTTA GCATGTACAC TTTTAGGTGT ACTTTCTGAA
701 GCCATGGGTTTAGAATTAGA GGCACCTACT AAAGCTTG TGATATGGA
751 CCAAAGATTGTGGTTAATT ACTACCCTAA GTGCCCTCAA CCTGACCTTA
801 CTTTAGGGCTCAAGAGGCAC ACCGACCCCG GGACTATAAC CCTCCTCCTT
851 CAGGACCAAGTCGGCGGTCT TCAGGCCACT CGTGACGGTG GTAAACTTG
901 GATTACCGTG CAGCCGGTTC CCGGTGCCTT CGTTGTAAAC CTTGGTGATC
951 ATGGTCATGTTGCGCGAAAA ATGGCCAAAG ACCTTGAGAT CGCCCGTCAT
1001 AAGAGGCTTG CTAAAGAGGA AATGCCTTTT AAAGAGTTGG ACGAGGCCAA
1051 GTTTGAGTCC AAATCTATTG ACCAAATACT TGCTTAGATG GGCTTGGTTT
1101 GGTTCATTA TATTAAATTT ATTATTATTA TTATTTATTG CATTTGATAT
1151 GATATGATTG GAAATAAAAG AGAGATTGTT TGTGATAATT TGTGTGATTA
1201 TTATATCACT AAGTTATGGC TTTAATTGT GGTATGTTGG GAATTATATA
1251 TTTAGTTTTG TGTGAAGAAT ATATGATTTA AAGTTAAAAA AAAAAATGAT
1301 TTGTTATATGATTTACTTGT AAGGTTATAA GGTTATATTT ATTGTTGAG
1351 TTTGCGTATA

Fig. 12

12/18




Fig. 11

15/18




Fig. 14

14/18

1 CCAAGCCCAT CTAAGCAAGT ATTGGTCAA TAGATTGGA CTCAAACTTG GCCTCGTCCA
61 ACTCTTTAAA AGGCATTTC TCCTTAGCAA GCCTCTTATG ACGGGCGATC TCAAGGTCTT
121 TGGCCATTTT TCGCCGAACA TGACCATGAT CACCAAGGTT AACAAAGAA GCAACGGGAA
181 CCGGCTGCAC GGTAATCCAA GTTTACCAC CGTCACGAGT GGCCTGAAGA CCGCCGACTT
241 GGTCTGAAG GAGGAGGTT ATAGTCCCG GGTCCGTGTG CCTCTTGAGC CCTAAAGTAA
301 GGTACGGTTG AGGGCACTTA GGGTAGTAAT TAACCACAAT CTTTGGTCC ATATCAACAC
361 AAGCTTTAGT AAGTGCCCTCT AATTCTAAAC CCATGGCTTC AGAAAGTACA CCTAAAAGTG
421 TACATGCTAA GGTCATTAAAC TTGTTGCTGT ATTCTCTGT GACCTTTATC CAACCCCTCTG
481 GTTTGCTCG CCATCTTGTG TAGTCCCTTG AGTTCGTGCG GTATGAGAAA TACGTACCGA
541 TTCCCTCCA GTCCTGCACT ACTTCTCCCT GTTGATTGAA CCTGAAGATG ACTCGACACG
601 ATAAACCCGC CCTTTTACC ACCAGACATA TCAAATCGGA GCTTCTCTTC TGCCGGGAGA
661 GCGAAAAATT CACGAGCCAA CCGAGTCATA TCAGCAATAA GATCGTCACC AACACCGTGA
721 TCAACCACTT GAAAAATCCC CCAATCTTCA CACGCCTCAA CAATCTTCCG ACATATTICA
781 CCCCTTTTTT CACCATCTAT ACCAGCAAGA GATATCACCG GAATATCATT GCTAAACTCA
841 TTATACGCCA CTTTCGGAGC TTCATCCTCG TCCCTAACAA AATTCGAGTT CAATTTATCG
901 TCCCCTTCTA GTGAAGTGAG CGTTTGGG

Fig. 13

17/18




Fig. 16

16/18




Fig. 15

18/18




Fig. 17



— Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(88) Date of publication of the international search report:
14 December 2000

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
31 August 2000 (31.08.2000)


PCT

(10) International Publication Number
WO 00/50613 A3

- (51) International Patent Classification⁷: C12N 15/82, 74040 Nes Ziona (IL). OVADIS, Marianna [IL/IL];
A01H 5/00 Hershenson Street 76/28, 76484 Rehovot (IL).
- (21) International Application Number: PCT/IL00/00110 (74) Agent: REINHOLD COHN AND PARTNERS; P.O.
Box 4060, 61040 Tel Aviv (IL).
- (22) International Filing Date: 22 February 2000 (22.02.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/121,239 22 February 1999 (22.02.1999) US
- (71) Applicant (for all designated States except US): YISSUM RESEARCH AND DEVELOPMENT COMPANY OF THE HEBREW UNIVERSITY OF JERUSALEM [IL/IL]; Jabotinsky Street 46, 91042 Jerusalem (IL).
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): VAINSTEIN, Alexander [IL/IL]; Hanassi Harishron Street 42/6, 76303 Rehovot (IL). ZUKER, Amir [IL/IL]; Negba Street 32/18,
- Published: — With international search report.

[Continued on next page]

(54) Title: TRANSGENIC PLANTS AND METHOD FOR TRANSFORMING CARNATIONS



(57) Abstract: A method of transforming a carnation (*Dianthus L.*) plant genome with a DNA molecule. The method comprises (a) preparing stem explants from carnation cuttings; (b) wounding the explants by microprojectile bombardment; (c) cocultivating the wounded explants with *Agrobacterium* comprising the DNA molecule under defined conditions of exposure to dark followed by light; (d) excising shoots from the cultivated wounded explants and removing the leaves from the shoots; and (e) culturing the leaves to obtain transgenic shoots transformed with the DNA molecule. Also disclosed are a *rolC*-transgenic carnation with improved agronomic traits and enhancement of flower fragrance by antisense suppression of the flavonoid gene *fla*.

WO 00/50613 A3

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IL 00/00110

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	OVADIS, M., ET AL.: "Generation of transgenic carnation plants with novel characteristics by combining microprojectile bombardment with Agrobacterium tumefaciens transformation" CURRENT PLANT SCIENCE AND BIOTECHNOLOGY IN AGRICULTURE VOL. 36., pages 189-192, XP000921439 Plant biotechnology and in vitro biology in the 21st century. Proceedings of the IXth International Congress of the International Association of Plant Tissue Culture and Biotechnology, Jerusalem, Israel, 14-19 June 1998. ISBN: 0-7923-5826-0. 'EDITOR!; Ziv, M. 'EDITOR!; Izhar, S. 'EDITOR! the whole document	1-9
Y	& MSC. THESIS. AHRONI A.: 1996 (THE HEBREW UNIVERSITY OF JERUSALEM ISRAEL),	1-9
Y	ZUKER, A., ET AL.: "A highly efficient method for carnation transformation" ACTA HORTICULTURAE, (1997) NO. 447, PP. 373-375., XP000921438 Proceedings of the third international ISHS symposium on in vitro culture and horticultural breeding, Jerusalem, Israel, 16-21 June, 1996. ISSN: 0567-7572; ISBN: 90-6605-909-5. the whole document	1-9
Y	WO 92 17056 A (FLORIGENE B V) 15 October 1992 (1992-10-15)	1-9
A	the whole document	21-28
Y	FIROOZABADY, E., ET AL.: "Efficient transformation and regeneration of carnation cultivars using Agrobacterium" MOLECULAR BREEDING, vol. 1, 1995, pages 283-293, XP002140940 the whole document	1-9
X	TANAKA YOSHIKAZU ET AL: "Metabolic engineering to modify flower color." PLANT AND CELL PHYSIOLOGY, vol. 39, no. 11, November 1998 (1998-11), pages 1119-1126, XP000922977 ISSN: 0032-0781 page 1122, right-hand column -page 1125 & ZUKER, A., ET AL.: ABST. IX. INTERNATIONAL CONGRESS ON PLANT TISSUE AND CELL CULTURE, 1998, page 35	14-20, 26-28

-/--

INTERNATIONAL SEARCH REPORT

In tional Application No

PCT/IL 00/00110

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C12N15/82 A01H5/00		
According to International Patent Classification (IPC) or to both national classification and IFC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 C12N A01H		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, BIOSIS, WPI Data, CAB Data, EMBL		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	OVADIS, M., ET AL.: "A highly efficient procedure for generating carnation plants with novel traits" PROCEEDINGS OF THE NINETEENTH INTERNATIONAL SYMPOSIUM ON IMPROVEMENT OF ORNAMENTAL PLANTS. BREEDING ORNAMENTALS IN THE FUTURE: GOALS, GENES, TOOLS, ANGERS, FRANCE, 27-30 JULY, 1998. ACTA HORTICULTURAE, (2000) NO. 508, PP. 49-51., XP000946845 ISSN: 0567-7572; ISBN: 90-6605-832-3 the whole document --- -/-	1-9, 14-20, 26-28
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C.		
<input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : <div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search 6 October 2000		Date of mailing of the international search report 18.10.00
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer Maddox, A

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IL 00/00110

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GUTTERSON NEAL: "Anthocyanin biosynthetic genes and their application to flower color modification through sense suppression." HORTSCIENCE, vol. 30, no. 5, 1995, pages 964-966, XP000922821 ISSN: 0018-5345 the whole document	26-28
X	WO 96 36716 A (INT FLOWER DEV PTY LTD ;HOLTON TIMOTHY ALBERT (AU)) 21 November 1996 (1996-11-21) the whole document	26-28
P,X	ZUKER AMIR ET AL: "Wounding by bombardment yields highly efficient Agrobacterium-mediated transformation of carnation (Dianthus caryophyllus L.):" MOLECULAR BREEDING, vol. 5, no. 4, 1999, pages 367-375, XP002140941 ISSN: 1380-3743 the whole document	1-9
A	WO 94 28140 A (HOLTON TIMOTHY ALBERT ;INT FLOWER DEV PTY LTD (AU); TANAKA YOSHIKA) 8 December 1994 (1994-12-08) example 24	1-9
A	ZUKER AMIR ET AL: "Transformation of carnation by microprojectile bombardment." SCIENTIA HORTICULTURAE (AMSTERDAM), vol. 64, no. 3, 1995, pages 177-185, XP000921441 ISSN: 0304-4238 the whole document	1-9
A	WATAD ABED A ET AL: "Adventitious shoot formation from carnation stem segments: A comparison of different culture procedures." SCIENTIA HORTICULTURAE (AMSTERDAM), vol. 65, no. 4, 1996, pages 313-320, XP000921442 ISSN: 0304-4238 the whole document	1-9
A	WO 95 06741 A (BIOCEM ;KNITTEL NATHALIE (AU); LENEÉ PHILIPPE (FR)) 9 March 1995 (1995-03-09) the whole document	1-9

-/-

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IL 00/00110

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DEDIO J ET AL: "Molecular cloning of the flavanone 3-beta-hydroxylase gene (FHT) from carnation (<i>Dianthus caryophyllus</i>) and analysis of stable and unstable FHT mutants."</p> <p>THEORETICAL AND APPLIED GENETICS, vol. 90, no. 5, 1995, pages 611-617, XP000915009</p> <p>ISSN: 0040-5752</p> <p>the whole document</p> <p>-& DATABASE EMBL 'Online!</p> <p>ACCESSION NO: X70378,</p> <p>23 February 1993 (1993-02-23)</p> <p>DEDIO, J., ET AL.: "D.caryophyllus fht gene for narigenin 3-dioxygenase"</p> <p>XP002149237</p> <p>the whole document</p>	14,18, 26-28
X	<p>BRITSCH L ET AL: "Molecular characterization of flavone-3'-beta'-hydroxylases"</p> <p>EUROPEAN JOURNAL OF BIOCHEMISTRY, DE, BERLIN, vol. 217, no. 2, 15 October 1993 (1993-10-15), pages 745-754, XP002108555</p> <p>ISSN: 0014-2956</p> <p>the whole document</p> <p>-& DATABASE EMBL 'Online!</p> <p>ACCESSION NO: X72592,</p> <p>16 March 1993 (1993-03-16)</p> <p>BRITSCH L., ET AL.: "D.caryophyllus mRNA for flavanone 3-beta-hydroxylase"</p> <p>XP002149238</p> <p>the whole document</p>	20
X	<p>WO 96 39827 A (PROGUARD INC ;EMERSON RALPH W (US); CRANDALL BRADFORD G JR (US))</p> <p>19 December 1996 (1996-12-19)</p> <p>page 5, line 1 - line 5; example 11</p> <p>page 12, line 20 -page 19, line 30</p>	21,23,26
X	<p>WO 96 20595 A (PROGUARD INC ;EMERSON RALPH W (US); CRANDALL BRADFORD G JR (US))</p> <p>11 July 1996 (1996-07-11)</p> <p>page 4, line 25 - line 28</p> <p>page 16, line 31 -page 23, line 17;</p> <p>examples 8,9</p>	21-23,26
X	<p>WO 97 35471 A (PROGUARD INC ;EMERSON RALPH W (US); CRANDALL BRADFORD G JR (US))</p> <p>2 October 1997 (1997-10-02)</p> <p>page 7, line 7 - line 11</p> <p>page 19, line 3 -page 32, line 10; example 31</p>	21,23,26

-/-

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-9

Method for transforming carnation comprising a combined microprojectile/Agrobacterium protocol.

2. Claims: 10-13

Transgenic carnations transformed with the rolC gene

3. Claims: 14-28

Methods for controlling fragrance/colour of plants by modulating the anthocanin-biosynthetic pathway, specifically modulating the FHT gene expression, transgenic carnations with modulated FHT expression, and the DNA sequence of fig.13

INTERNATIONAL SEARCH REPORT

International Application No
PCT/IL 00/00110

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 486 233 A (PIONEER HI BRED INT) 20 May 1992 (1992-05-20) the whole document	1-9
A	WO 93 18142 A (UNIV WASHINGTON) 16 September 1993 (1993-09-16) page 5 -page 11	14-28
A	WO 97 15584 A (UNIV MICHIGAN) 1 May 1997 (1997-05-01) page 15, line 7 - line 21	21-26
A	ZUKER A ET AL: "Genetic engineering for cut-flower improvement" BIOTECHNOLOGY ADVANCES,GB,ELSEVIER PUBLISHING, BARKING, vol. 16, no. 1, 1998, pages 33-79, XP004099136 ISSN: 0734-9750 page-58, paragraph 3	21-26
A	PELLEGRINESCHI A ET AL: "IMPROVEMENT OF ORNAMENTAL CHARACTERS AND FRAGRANCE PRODUCTION IN LEMON-SCENTED GERANIUM THROUGH GENETIC TRANSFORMATION BY AGROBACTERIUM RHIZOGENES" BIO/TECHNOLOGY,US,NATURE PUBLISHING CO. NEW YORK, vol. 12, no. 1, 1994, pages 64-68, XP002036390 ISSN: 0733-222X the whole document	21-26
A	WO 98 50570 A (UNIV MICHIGAN ;WANG JIHONG (US); PICHESKY ERAN (US)) 12 November 1998 (1998-11-12) the whole document	21-26
A	WO 97 21816 A (ZENECA LTD ;MANNING KENNETH (GB)) 19 June 1997 (1997-06-19) page 6, line 16 - line 18; table 1	21-26

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IL 00/00110

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0486233 A		DK 486233 T	11-09-1995
		ES 2077181 T	16-11-1995
		GR 3017619 T	31-01-1996
		HU 60783 A	28-10-1992
		JP 5308960 A	22-11-1993
		NZ 239977 A	26-08-1993
WO 9318142 A	16-09-1993	AU 668548 B	09-05-1996
		AU 3786493 A	05-10-1993
		BG 99033 A	29-09-1995
		BR 9306047 A	18-11-1997
		CA 2131704 A	16-09-1993
		EP 0630403 A	28-12-1994
		HU 69991 A	28-09-1995
		JP 7506963 T	03-08-1995
		NZ 251069 A	26-01-1996
		US 5733759 A	31-03-1998
WO 9715584 A	01-05-1997	AU 7663296 A	15-05-1997
		EP 0858507 A	19-08-1998
WO 9850570 A	12-11-1998	AU 7377298 A	27-11-1998
WO 9721816 A	19-06-1997	AU 1106297 A	03-07-1997

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IL 00/00110

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9217056 A	15-10-1992	EP 0582603 A US 5589613 A	16-02-1994 31-12-1996
WO 9639827 A	19-12-1996	AU 4648996 A BR 9508561 A CA 2196972 A EP 0776159 A NZ 301140 A US 5792467 A ZA 9600006 A	30-12-1996 19-05-1998 19-12-1996 04-06-1997 23-12-1998 11-08-1998 27-08-1996
WO 9620595 A	11-07-1996	AU 4744296 A BR 9510180 A CA 2208753 A EP 0800345 A EP 1000543 A JP 11501295 T PL 321179 A ZA 9600005 A	24-07-1996 14-10-1997 11-07-1996 15-10-1997 17-05-2000 02-02-1999 24-11-1997 27-08-1996
WO 9735471 A	02-10-1997	AU 3202597 A CA 2250182 A EP 0889691 A	17-10-1997 02-10-1997 13-01-1999
WO 9636716 A	21-11-1996	AU 699874 B AU 5639696 A CA 2202668 A EP 0873410 A HU 9802555 A JP 11505116 T NZ 307119 A US 6080920 A	17-12-1998 29-11-1996 21-11-1996 28-10-1998 01-02-1999 18-05-1999 29-04-1999 27-06-2000
WO 9428140 A	08-12-1994	AU 672308 B AU 6789594 A CA 2163220 A CN 1127015 A EP 0703982 A JP 8511683 T NZ 266401 A PL 311691 A SG 45175 A US 5948955 A	26-09-1996 20-12-1994 08-12-1994 17-07-1996 03-04-1996 10-12-1996 25-09-1996 04-03-1996 16-01-1998 07-09-1999
WO 9506741 A	09-03-1995	FR 2709496 A AU 695398 B AU 7539694 A EP 0716706 A HU 74664 A JP 9509302 T SK 25896 A	10-03-1995 13-08-1998 22-03-1995 19-06-1996 28-01-1997 22-09-1997 04-09-1996
EP 0486233 A	20-05-1992	AT 125301 T AU 643846 B AU 8771391 A CA 2053473 A DE 69111385 D DE 69111385 T	15-08-1995 25-11-1993 21-05-1992 15-05-1992 24-08-1995 04-01-1996

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IL 00/00110

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/82 A01H5/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, WPI Data, CAB Data, EMBL

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	OVADIS, M., ET AL.: "A highly efficient procedure for generating carnation plants with novel traits" PROCEEDINGS OF THE NINETEENTH INTERNATIONAL SYMPOSIUM ON IMPROVEMENT OF ORNAMENTAL PLANTS. BREEDING ORNAMENTALS IN THE FUTURE: GOALS, GENES, TOOLS, ANGERS, FRANCE, 27-30 JULY, 1998. ACTA HORTICULTURAE, (2000) NO. 508, PP. 49-51., XP000946845 ISSN: 0567-7572; ISBN: 90-6605-832-3 the whole document --- -/--	1-9, 14-20, 26-28

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

23 October 2000

Date of mailing of the international search report

25.10.00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Maddox, A

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

REVISED VERSION

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
31 August 2000 (31.08.2000)

PCT

(10) International Publication Number
WO 00/50613 A3

(51) International Patent Classification⁷: C12N 15/82, A01H 5/00

(21) International Application Number: PCT/IL00/00110

(22) International Filing Date: 22 February 2000 (22.02.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/121,239 22 February 1999 (22.02.1999) US

(71) Applicant (for all designated States except US): YISSUM RESEARCH AND DEVELOPMENT COMPANY OF THE HEBREW UNIVERSITY OF JERUSALEM [IL/IL]; Jabotinsky Street 46, 91042 Jerusalem (IL).

(72) Inventors; and

(75) Inventors/Applicants (for US only): VAINSTEIN, Alexander [IL/IL]; Hanassi Harishron Street 42/6, 76303 Rehovot (IL). ZUKER, Amir [IL/IL]; Negba Street 32/18, 74040 Nes Ziona (IL). OVADIS, Marianna [IL/IL]; Hershenon Street 76/28, 76484 Rehovot (IL).

(74) Agent: REINHOLD COHN AND PARTNERS; P.O. Box 4060, 61040 Tel Aviv (IL).

(81) Designated States (national): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— With international search report.


(88) Date of publication of the international search report:
14 December 2000

Date of publication of the revised international search report:
22 February 2001

(15) Information about Correction:
see PCT Gazette No. 08/2001 of 22 February 2001, Section II

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: TRANSGENIC PLANTS AND METHOD FOR TRANSFORMING CARNATIONS



(57) Abstract: A method of transforming a carnation (*Dianthus L.*) plant genome with a DNA molecule. The method comprises (a) preparing stem explants from carnation cuttings; (b) wounding the explants by microprojectile bombardment; (c) cocultivating the wounded explants with *Agrobacterium* comprising the DNA molecule under defined conditions of exposure to dark followed by light; (d) excising shoots from the cultivated wounded explants and removing the leaves from the shoots; and (e) culturing the leaves to obtain transgenic shoots transformed with the DNA molecule. Also disclosed are a *rolC*-transgenic carnation with improved agronomic traits and enhancement of flower fragrance by antisense suppression of the flavonoid gene *fht*.



WO 00/50613 A3

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IL 00/00110

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DEDIO J ET AL: "Molecular cloning of the flavanone 3-beta-hydroxylase gene (FHT) from carnation (<i>Dianthus caryophyllus</i>) and analysis of stable and unstable FHT mutants."</p> <p>THEORETICAL AND APPLIED GENETICS, vol. 90, no. 5, 1995, pages 611-617, XP000915009</p> <p>ISSN: 0040-5752</p> <p>the whole document</p> <p>-& DATABASE EMBL [Online]</p> <p>ACCESSION NO: X70378,</p> <p>23 February 1993 (1993-02-23)</p> <p>DEDIO, J., ET AL.: "D.caryophyllus fht gene for narigenin 3-dioxygenase"</p> <p>XP002149237</p> <p>the whole document</p>	14,18, 26-28
X	<p>BRITSCH L ET AL: "Molecular characterization of flavone-3[beta]-hydroxylases"</p> <p>EUROPEAN JOURNAL OF BIOCHEMISTRY, DE, BERLIN, vol. 217, no. 2, 15 October 1993 (1993-10-15), pages 745-754, XP002108555</p> <p>ISSN: 0014-2956</p> <p>the whole document</p> <p>-& DATABASE EMBL [Online]</p> <p>ACCESSION NO: X72592,</p> <p>16 March 1993 (1993-03-16)</p> <p>BRITSCH L., ET AL.: "D.caryophyllus mRNA for flavanone 3-beta-hydroxylase"</p> <p>XP002149238</p> <p>the whole document</p>	20
X	<p>WO 96 39827 A (PROGUARD INC ;EMERSON RALPH W (US); CRANDALL BRADFORD G JR (US))</p> <p>19 December 1996 (1996-12-19)</p> <p>page 5, line 1 - line 5; example 11</p> <p>page 12, line 20 -page 19, line 30</p>	21,23,26
X	<p>WO 96 20595 A (PROGUARD INC ;EMERSON RALPH W (US); CRANDALL BRADFORD G JR (US))</p> <p>11 July 1996 (1996-07-11)</p> <p>page 4, line 25 - line 28</p> <p>page 16, line 31 -page 23, line 17;</p> <p>examples 8,9</p>	21-23,26
X	<p>WO 97 35471 A (PROGUARD INC ;EMERSON RALPH W (US); CRANDALL BRADFORD G JR (US))</p> <p>2 October 1997 (1997-10-02)</p> <p>page 7, line 7 - line 11</p> <p>page 19, line 3 -page 32, line 10; example 31</p>	21,23,26

-/--

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IL 00/00110

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	OVADIS, M., ET AL.: "Generation of transgenic carnation plants with novel characteristics by combining microprojectile bombardment with Agrobacterium tumefaciens transformation" CURRENT PLANT SCIENCE AND BIOTECHNOLOGY IN AGRICULTURE VOL. 36., pages 189-192, XP000921439 Plant biotechnology and in vitro biology in the 21st century. Proceedings of the IXth International Congress of the International Association of Plant Tissue Culture and Biotechnology, Jerusalem, Israel, 14-19 June 1998. ISBN: 0-7923-5826-0. [EDITOR]; Ziv, M. [EDITOR]; Izhar, S. [EDITOR] the whole document	1-9
Y	& MSC THESIS. AHRONI A.. 1996 (THE HEBREW UNIVERSITY OF JERUSALEM ISRAEL),	1-9
Y	--- ZUKER, A., ET AL.: "A highly efficient method for carnation transformation" ACTA HORTICULTURAE, (1997) NO. 447, PP. 373-375., XP000921438 Proceedings of the third international ISHS symposium on in vitro culture and horticultural breeding, Jerusalem, Israel, 16-21 June, 1996. ISSN: 0567-7572; ISBN: 90-6605-909-5 the whole document	1-9
Y	--- WO 92 17056 A (FLORIGENE B V) 15 October 1992 (1992-10-15)	1-9
A	the whole document	21-28
Y	--- FIROOZABADY, E., ET AL.: "Efficient transformation and regeneration of carnation cultivars using Agrobacterium" MOLECULAR BREEDING, vol. 1, 1995, pages 283-293, XP002140940 the whole document	1-9
X	--- TANAKA YOSHIKAZU ET AL: "Metabolic engineering to modify flower color." PLANT AND CELL PHYSIOLOGY, vol. 39, no. 11, November 1998 (1998-11), pages 1119-1126, XP000922977 ISSN: 0032-0781 page 1122, right-hand column -page 1125 & ZUKER, A., ET AL.: ABST. IX. INTERNATIONAL CONGRESS ON PLANT TISSUE AND CELL CULTURE, 1998, page 35 --- -/--	14-20, 26-28

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IL 00/00110

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 95 06741 A (BIOCEM ;KNITTEL NATHALIE (AU); LENEÉ PHILIPPE (FR)) 9 March 1995 (1995-03-09) the whole document	1-9
A	EP 0 486 233 A (PIONEER HI BRED INT) 20 May 1992 (1992-05-20) the whole document	1-9
A	WO 93 18142 A (UNIV WASHINGTON) 16 September 1993 (1993-09-16) page 5 -page 11	14-28
A	WO 97 15584 A (UNIV MICHIGAN) 1 May 1997 (1997-05-01) page 15, line 7 - line 21	21-26
A	ZUKER A ET AL: "Genetic engineering for cut-flower improvement" BIOTECHNOLOGY ADVANCES,GB,ELSEVIER PUBLISHING, BARKING, vol. 16, no. 1, 1998, pages 33-79, XP004099136 ISSN: 0734-9750 page 58, paragraph 3	21-26
A	PELLEGRINESCHI A ET AL: "IMPROVEMENT OF ORNAMENTAL CHARACTERS AND FRAGRANCE PRODUCTION IN LEMON-SCENTED GERANIUM THROUGH GENETIC TRANSFORMATION BY AGROBACTERIUM RHIZOGENES" BIO/TECHNOLOGY,US,NATURE PUBLISHING CO. NEW YORK, vol. 12, no. 1, 1994, pages 64-68, XP002036390 ISSN: 0733-222X the whole document	21-26
A	WO 98 50570 A (UNIV MICHIGAN ;WANG JIHONG (US); PICHESKY ERAN (US)) 12 November 1998 (1998-11-12) the whole document	21-26
A	WO 97 21816 A (ZENECA LTD ;MANNING KENNETH (GB)) 19 June 1997 (1997-06-19) page 6, line 16 - line 18; table 1	21-26

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IL 00/00110

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GUTTERSON NEAL: "Anthocyanin biosynthetic genes and their application to flower color modification through sense suppression." HORTSCIENCE, vol. 30, no. 5, 1995, pages 964-966, XP000922821 ISSN: 0018-5345 the whole document ---	26-28
X	WO 96 36716 A (INT FLOWER DEV PTY LTD ;HOLTON TIMOTHY ALBERT (AU)) 21 November 1996 (1996-11-21) the whole document ---	26-28
P,X	ZUKER AMIR ET AL: "Wounding by bombardment yields highly efficient Agrobacterium-mediated transformation of carnation (Dianthus caryophyllus L.)." MOLECULAR BREEDING, vol. 5, no. 4, 1999, pages 367-375, XP002140941 ISSN: 1380-3743 the whole document ---	1-9
P,X	WO 99 37794 A (HUGHES STEPHEN GLYN ;VERHOEYEN MARTINE ELISA (GB); UNILEVER PLC (G) 29 July 1999 (1999-07-29) example 11 ---	21,26
A	WO 94 28140 A (HOLTON TIMOTHY ALBERT ;INT FLOWER DEV PTY LTD (AU); TANAKA YOSHIKA) 8 December 1994 (1994-12-08) example 24 ---	1-9
A	ZUKER AMIR ET AL: "Transformation of carnation by microprojectile bombardment." SCIENTIA HORTICULTURAE (AMSTERDAM), vol. 64, no. 3, 1995, pages 177-185, XP000921441 ISSN: 0304-4238 the whole document ---	1-9
A	WATAD ABED A ET AL: "Adventitious shoot formation from carnation stem segments: A comparison of different culture procedures." SCIENTIA HORTICULTURAE (AMSTERDAM), vol. 65, no. 4, 1996, pages 313-320, XP000921442 ISSN: 0304-4238 the whole document ---	1-9

-/--

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-9

Method for transforming carnation comprising a combined microprojectile/Agrobacterium protocol.

2. Claims: 10-13

Transgenic carnations transformed with the rolC gene

3. Claims: 14-28

Methods for controlling fragrance/colour of plants by modulating the anthocanin-biosynthetic pathway, specifically modulating the FHT gene expression, transgenic carnations with modulated FHT expression, and the DNA sequence of fig.13

INTERNATIONAL SEARCH REPORT

International application No.
PCT/IL 00/00110

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☒ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
1-9(subject group 1) and 14-28(subject group 3)

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.